

# The **mandi** Bundle

Paul J. Heafner\*

January 26, 2022

mandi version v3.1.0 dated 2022-01-27  
mandistudent version v3.1.0 dated 2022-01-27  
mandiexp version v3.1.0 dated 2022-01-27

---

\*Email: [heafnerj@gmail.com](mailto:heafnerj@gmail.com)

# Contents

<b>Acknowledgements</b>	<b>3</b>
<b>Change History</b>	<b>4</b>
<b>List of Web VPython Programs</b>	<b>5</b>
<b>List of VPython Programs</b>	<b>5</b>
<b>List of Figures</b>	<b>5</b>
<b>1 Introduction</b>	<b>6</b>
<b>2 Code Availability</b>	<b>6</b>
<b>3 Student/Instructor Quick Guide</b>	<b>7</b>
<b>4 The <code>mandi</code> Package</b>	<b>8</b>
4.1 Package Options . . . . .	8
4.2 The <code>mandisetup</code> Command . . . . .	8
4.3 Lua <sup>L</sup> TeX is Required . . . . .	8
4.4 Physical Quantities . . . . .	9
4.4.1 Typesetting Physical Quantities . . . . .	9
4.4.2 Checking Physical Quantities . . . . .	10
4.4.3 Predefined Physical Quantities . . . . .	10
4.4.4 Defining and Redefining Physical Quantities . . . . .	22
4.4.5 Changing Units . . . . .	22
4.5 Physical Constants . . . . .	23
4.5.1 Typesetting Physical Constants . . . . .	23
4.5.2 Checking Physical Constants . . . . .	24
4.5.3 Predefined Physical Constants . . . . .	24
4.5.4 Defining and Redefining Physical Constants . . . . .	30
4.5.5 Changing Precision . . . . .	31
4.6 Predefined Units and Constructs . . . . .	31
4.7 <code>mandi</code> Source Code . . . . .	35
<b>5 The <code>mandistudent</code> Package</b>	<b>51</b>
5.1 Traditional Vector Notation . . . . .	51
5.2 Problems and Annotated Problem Solutions . . . . .	55
5.3 Coordinate-Free and Index Notation . . . . .	60
5.4 Web VPython and VPython Program Listings . . . . .	62
5.5 The <code>webvpythonblock</code> Environment . . . . .	62
5.6 The <code>vpythonfile</code> Command . . . . .	68
5.7 The <code>webvpythoninline</code> and <code>vpythoninline</code> Commands . . . . .	70
5.8 <code>mandistudent</code> Source Code . . . . .	71
<b>6 The <code>mandiexp</code> Package</b>	<b>82</b>
6.1 The Fundamental Principles . . . . .	82
6.2 Other Expressions . . . . .	84
6.3 <code>mandiexp</code> Source Code . . . . .	87
<b>7 Index</b>	<b>93</b>

## Acknowledgements

To all of the students who have learned L<sup>A</sup>T<sub>E</sub>X 2 <sub>$\varepsilon$</sub>  in my introductory physics courses over the years, I say a heartfelt thank you. You have contributed directly to the state of this software and to its use in introductory physics courses and to innovating how physics is taught.

I also acknowledge the L<sup>A</sup>T<sub>E</sub>X 2 <sub>$\varepsilon$</sub>  developers who inhabit the [TeX StackExchange](#) site. Entering a new culture is daunting for anyone, especially for newcomers; the L<sup>A</sup>T<sub>E</sub>X 2 <sub>$\varepsilon$</sub>  development culture is no exception. We all share a passion for creating beautiful documents and I have learned much over the summers of 2020 and 2021 that improved my ability to do just that. There are too many of you to list individually, and I would surely accidentally omit some were I to try. Collectively, I thank you all for your patience and advice.

# Change History

v3.0.0 (2021-08-22)	
<code>mandiexp</code> → P.82 Initial release	6
<code>mandi</code> → P.8 Initial release	6
<code>mandistudent</code> → P.51 Initial release	6
v3.0.1 (2021-08-24)	
<code>mandiexp</code> → P.82 Minor doc changes	6
<code>mandi</code> → P.8 Minor doc changes	6
<code>mandistudent</code> → P.51 Minor doc changes	6
v3.1.0 (2022-01-27)	
<code>mandiexp</code> → P.82 Code formatted for better readability	87
<code>mandiexp</code> → P.82 Version number works	87
<code>mandiexp</code> → P.82 <code>xparse</code> is loaded for older formats	87
<code>mandi</code> → P.8 Added GitHub links to code	6
<code>mandi</code> → P.8 Added <code>\hbar</code> → P.33	45
<code>mandi</code> → P.8 Added <code>\lorentzfactor</code> → P.16	43
<code>mandi</code> → P.8 Added a negative space to <code>\lightspeed</code> → P.32	37
<code>mandi</code> → P.8 Code formatted for better readability	35
<code>mandi</code> → P.8 Constants' values now use only <code>\times</code>	46
<code>mandi</code> → P.8 Improved <code>\checkconstant</code> → P.24	49
<code>mandi</code> → P.8 Improved <code>\checkquantity</code> → P.10	49
<code>mandi</code> → P.8 Unknown package options handled safely	36
<code>mandi</code> → P.8 <code>LATEX3</code> code now conforms to formatting standards	38
<code>mandi</code> → P.8 <code>\mivector</code> → P.34 now requires more than one component	50
<code>mandi</code> → P.8 <code>xparse</code> is loaded for older formats	35
<code>mandistudent</code> → P.51 All instances of <code>GlowScript</code> have been changed to Web VPython	61
<code>mandistudent</code> → P.51 Code formatted for better readability	71
<code>mandistudent</code> → P.51 Default URL for <code>\vpythonfile</code> → P.68 is now <code>vpython.org</code>	80
<code>mandistudent</code> → P.51 Slightly modified <code>\image</code> → P.59	76
<code>mandistudent</code> → P.51 URLs fixed in <code>\vpythonfile</code> → P.68	80
<code>mandistudent</code> → P.51 URLs fixed in <code>webpythonblock</code> → P.62	80
<code>mandistudent</code> → P.51 Version number works	71
<code>mandistudent</code> → P.51 <code>LATEX3</code> code now conforms to formatting standards	75
<code>mandistudent</code> → P.51 <code>\diff</code> renamed to <code>\df</code> → P.61 for compatibility with the <code>numerica</code> package	61
<code>mandistudent</code> → P.51 <code>\dirvec</code> → P.52 no longer adds <code>\scriptspace</code> when no sub/superscript is given	74
<code>mandistudent</code> → P.51 <code>\vec</code> → P.51 no longer adds <code>\scriptspace</code> when no sub/superscript is given	74
<code>mandistudent</code> → P.51 <code>webpythonblock*</code> → P.62 is a variant of <code>webpythonblock</code> → P.62 that omits the QR code	80
<code>mandistudent</code> → P.51 <code>webpythonblock</code> → P.62 now automatically generates QR codes for program listings	80
<code>mandistudent</code> → P.51 <code>xparse</code> is loaded for older formats	71

## **List of Web VPython Programs**

1	A Web VPython Program With QR Code . . . . .	64
2	A Web VPython Program Without QR Code . . . . .	67

## **List of VPython Programs**

1	A VPython Program . . . . .	69
---	-----------------------------	----

## **List of Figures**

1	Image shown 20 percent actual size. . . . .	59
2	Image shown 20 percent actual size and rotated. . . . .	60

# 1 Introduction

The `mandi`<sup>1</sup> bundle consists of three packages: `mandi`, `mandistudent`, and `mandiexp`. Package `mandi`<sup>→ P.8</sup> provides the core functionality, namely correctly typesetting physical quantities and constants with their correct SI units as either scalars or vectors, depending on which is appropriate. Package `mandistudent`<sup>→ P.51</sup> provides other typesetting capability appropriate for written problem solutions. Finally, package `mandiexp`<sup>→ P.82</sup> provides commands for typesetting expressions from *Matter & Interactions*<sup>2</sup>.

`mandi` has been completely rewritten from the ground up. It had gotten too large and clumsy to use and maintain. It (unknowingly) used deprecated packages. It had too many arcane “features” that were never used. It did not support Unicode. It was not compatible with modern engines, like Lua<sup>LATEX</sup>. It did not have a key-value interface. Options could not be changed on the fly within a document. In short, it was a mess. I hope this rewrite addresses all of the bad things and forms a better code base for maintenance, useability, and future improvements.

So many changes have been made that I think the best approach for former, as well as new, users is to treat this as a brand new experience. I think the most important thing to keep in mind is that I assume users, especially new users, will have a relatively recent TeX distribution (like TeX Live) that includes a recently updated L<sup>A</sup>T<sub>E</sub>X 2 <sub>$\varepsilon$</sub>  kernel. If users report that this is a major problem, I can provide some degree of backwards compatibility. However, I use a fully updated TeX Live distribution.

# 2 Code Availability

The `mandi` source repository’s `main` branch is at <https://github.com/heafnerj/mandi>. This code will usually coincide with that found on CTAN. The very latest build can be found on the `dev` branch found at <https://github.com/heafnerj/mandi/tree/dev>. Students and other academic academic users should probably get the `dev` branch code since it is stable and may contain improvements over the `main` branch code.

---

<sup>1</sup>The bundle name can be pronounced either with two syllables, to rhyme with *candy*, or with three syllables, as *M and I*.

<sup>2</sup>See *Matter & Interactions* and <https://matterandinteractions.org/> for details.

### 3 Student/Instructor Quick Guide

Use `\vecP.51` to typeset the symbol for a vector. Use `\magnitudeP.54` to typeset the symbol for a vector's magnitude. Use `\dirvecP.52` to typeset the symbol for a vector's direction. Use `\changeinP.52` to typeset the symbol for the change in a vector or scalar. Use `\zerovecP.52` to typeset the zero vector. Use `\timestentoP.34` to typeset scientific notation.

<code>\(\vec{p}\)</code> or <code>\(\vec{*p}\)</code>	$p$ or $\vec{p}$
<code>\(\vec{p}_{\text{final}}\)</code> or <code>\(\vec{*p}_{\text{final}}\)</code>	$p_{\text{final}}$ or $\vec{p}_{\text{final}}$
<code>\(\magnitude{\vec{p}}\)</code> or <code>\(\magnitude{\vec{*p}}\)</code>	$\ p\ $ or $\ p_{\text{final}}\ $
<code>\(\dirvec{p}\)</code> or <code>\(\dirvec{*p}\)</code>	$\hat{p}$ or $\hat{\vec{p}}$
<code>\(\changein{\vec{p}}\)</code> or <code>\(\changein{t}\)</code>	$\Delta p$ or $\Delta t$
<code>\(\zerovec\)</code> or <code>\(\zerovec{*}\)</code>	$\mathbf{0}$ or $\vec{0}$
<code>\(6.02\timestento{-19}\)</code>	$6.02 \times 10^{-19}$

Use a `physical quantity'sP.9` name to typeset a magnitude and that quantity's units. If the quantity is a vector, you can add `vector` either to the beginning or the end of the quantity's name. For example, if you want momentum, use `\momentumP.9` and its variants.

<code>\(\momentum{7.071}\)</code>	$7.071 \text{ kg} \cdot \text{m/s}$
<code>\(\vectormomentum{3,-4,5}\)</code>	$\langle 3, -4, 5 \rangle \text{ kg} \cdot \text{m/s}$
<code>\(\momentumvector{3,-4,5}\)</code>	$\langle 3, -4, 5 \rangle \text{ kg} \cdot \text{m/s}$

Use a `physical constant'sP.23` name to typeset its numerical value and units. Append `mathsymbol` to the constant's name to get its mathematical symbol. For example, if you want to typeset the vacuum permittivity, use `\vacuumpermittivityP.30` and its variant.

<code>\(\vacuumpermittivitymathsymbol = \vacuumpermittivity\)</code>	$\epsilon_0 = 9 \times 10^{-12} \text{ C}^2/\text{N} \cdot \text{m}^2$
--	--

Use `\mivectorP.34` to typeset symbolic vectors with components. Use the alias `\directionP.13` to typeset a direction or unit vector.

<code>\(\mivector{\slot,\slot,\slot}\)</code> or <code>\(\mivector{p_x,p_y,p_z}\)</code>	$\langle p_x, p_y, p_z \rangle$
--	---------------------------------

$\langle \frac{1}{\sqrt{3}}, \frac{1}{\sqrt{3}}, \frac{1}{\sqrt{3}} \rangle$ or $\langle p_x, p_y, p_z \rangle$
---

Use `\physicsproblemP.55` and `\partsP.55` and `\problempartP.55` for problems. For step-by-step mathematical solutions use `\physicssolutionP.56`. Use `\webpythonblockP.62` to typeset `Web VPython` programs. Use `\vpythonfileP.68` to typeset `VPython` program files.

## 4 The **mandi** Package

Load **mandi** as you would any package in your preamble.

```
\usepackage[options]{mandi}
```

### \mandiversion

Typesets the current version and build date.

```
The version is \mandiversion\ and is a stable build.
```

```
The version is v3.1.0 dated 2022-01-27 and is a stable build.
```

### 4.1 Package Options

N 2021-01-30 `units=<type of unit>` (initially unspecified, set to **alternate**)  
N 2021-01-30 `preciseconstants=<boolean>` (initially unspecified, set to **false**)

Now **mandi** uses a key-value interface for options. The `units` key can be set to **base**, **derived**, or **alternate**. The `preciseconstants` key is always either **true** or **false**.

### 4.2 The **mandisetup** Command

N 2021-02-17 `\mandisetup{(options)}`

Command to set package options on the fly after loadtime. This can be done in the preamble or inside the `\begin{document}... \end{document}` environment.

```
\mandisetup{units=base}
```

```
\mandisetup{preciseconstants}
```

```
\mandisetup{preciseconstants = false}
```

### 4.3 Lua<sup>A</sup>T<sub>E</sub>X is Required

In order to make use of better fonts and Unicode features, **mandi** now requires the Lua<sup>A</sup>T<sub>E</sub>X engine for processing documents. It will not work with other engines.

#### 4.4 Physical Quantities

#### 4.4.1 Typesetting Physical Quantities

Typesetting physical quantities and constants using semantically appropriate names, along with the correct [SI units](#), is the core function of `mandi`. Take momentum as the prototypical physical quantity in an introductory physics course.

```
\momentum{\langle magnitude\rangle}  
\momentumvector{\langle c_1, \dots, c_n \rangle}  
\vectormomentum{\langle c_1, \dots, c_n \rangle}
```

Command for momentum and its vector variants. The default units will depend on the options passed to `mandi` at load time. Alternate units are the default. Other units can be forced as demonstrated. The vector variants can take more than three components. Note the other variants for the quantity's value and units.

Commands that include the name of a physical quantity typeset units, so they shouldn't be used for algebraic or symbolic values of components. For example, one shouldn't use `\vectormomentum{mv_x,mv_y,mv_z}` but instead the generic `\mivector{mv_x,mv_y,mv_z}` instead.

## 4.4.2 Checking Physical Quantities

U 2022-01-27

`\checkquantity{<name>}`

Command to check and typeset the command, base units, derived units, and alternate units of a defined physical quantity.

## 4.4.3 Predefined Physical Quantities

Every other defined physical quantity can be treated similarly. Just replace `momentum` with the quantity's name. Obviously, the variants that begin with `\vector` will not be defined for scalar quantities. Here are all the physical quantities, with all their units, defined in `mandi`. Rememeber that units are not present with symbolic (algebraic) quantities, so do not use the `\vector` variants of these commands for symbolic components. Use `\mivectorP.34` instead.

`\acceleration{<magnitude>}`  
`\accelerationvector{<c1, ..., cn>}`  
`\vectoracceleration{<c1, ..., cn>}`

command	<code>\acceleration</code>	
base	<code>derived</code>	<code>alternate</code>
$m \cdot s^{-2}$	N/kg	$m/s^2$

`\amount{<magnitude>}`

command	<code>\amount</code>	
base	<code>derived</code>	<code>alternate</code>
mol	mol	mol

`\angularacceleration{<magnitude>}`  
`\angularaccelerationvector{<c1, ..., cn>}`  
`\vectorangularacceleration{<c1, ..., cn>}`

command	<code>\angularacceleration</code>	
base	<code>derived</code>	<code>alternate</code>
$rad \cdot s^{-2}$	$rad/s^2$	$rad/s^2$

`\angularfrequency{<magnitude>}`

command	<code>\angularfrequency</code>	
base	<code>derived</code>	<code>alternate</code>
$rad \cdot s^{-1}$	$rad/s$	$rad/s$

`\angularimpulse{<magnitude>}`  
`\angularimpulsevector{<c1, ..., cn>}`  
`\vectorangularimpulse{<c1, ..., cn>}`

command	<code>\angularimpulse</code>	
base	<code>derived</code>	<code>alternate</code>
$kg \cdot m^2 \cdot s^{-1}$	$kg \cdot m^2/s$	$kg \cdot m^2/s$

N 2021-02-24

```
\angularmomentum{\(magnitude)}
\angularmomentumvector{\(c_1, \dots, c_n)}
\vectorangularmomentum{\(c_1, \dots, c_n)}
```

command	\angularmomentum	
base	derived	alternate
$\text{kg} \cdot \text{m}^2 \cdot \text{s}^{-1}$	$\text{kg} \cdot \text{m}^2/\text{s}$	$\text{kg} \cdot \text{m}^2/\text{s}$

N 2021-02-24

```
\angularvelocity{\(magnitude)}
\angularvelocityvector{\(c_1, \dots, c_n)}
\vectorangularvelocity{\(c_1, \dots, c_n)}
```

command	\angularvelocity	
base	derived	alternate
$\text{rad} \cdot \text{s}^{-1}$	rad/s	rad/s

```
\area{\(magnitude)}
```

command	\area	
base	derived	alternate
$\text{m}^2$	$\text{m}^2$	$\text{m}^2$

```
\areachargedensity{\(magnitude)}
```

command	\areachargedensity	
base	derived	alternate
$\text{A} \cdot \text{s} \cdot \text{m}^{-2}$	$\text{C}/\text{m}^2$	$\text{C}/\text{m}^2$

```
\areamassdensity{\(magnitude)}
```

command	\areamassdensity	
base	derived	alternate
$\text{kg} \cdot \text{m}^{-2}$	$\text{kg}/\text{m}^2$	$\text{kg}/\text{m}^2$

```
\capacitance{\(magnitude)}
```

command	\capacitance	
base	derived	alternate
$\text{A}^2 \cdot \text{s}^4 \cdot \text{kg}^{-1} \cdot \text{m}^{-2}$	F	C/V

```
\charge{\(magnitude)}
```

command	\charge	
base	derived	alternate
A · s	C	C

N 2021-02-24

```
\cmagneticfield{\(magnitude)}
\cmagneticfieldvector{\(c_1, \dots, c_n\)}
\vectorcmagneticfield{\(c_1, \dots, c_n\)}
```

command	\cmagneticfield	
base	derived	alternate
$\text{kg} \cdot \text{m} \cdot \text{A}^{-1} \cdot \text{s}^{-3}$	N/C	N/C

```
\conductance{\(magnitude)}
```

command	\conductance	
base	derived	alternate
$\text{A}^2 \cdot \text{s}^3 \cdot \text{kg}^{-1} \cdot \text{m}^{-2}$	S	A/V

```
\conductivity{\(magnitude)}
```

command	\conductivity	
base	derived	alternate
$\text{A}^2 \cdot \text{s}^3 \cdot \text{kg}^{-1} \cdot \text{m}^{-3}$	S/m	A/V · m

```
\conventionalcurrent{\(magnitude)}
```

command	\conventionalcurrent	
base	derived	alternate
A	C/s	A

```
\current{\(magnitude)}
```

command	\current	
base	derived	alternate
A	A	A

```
\currentdensity{\(magnitude)}
```

```
\currentdensityvector{\(c_1, \dots, c_n\)}
\vectorcurrentdensity{\(c_1, \dots, c_n\)}
```

command	\currentdensity	
base	derived	alternate
$\text{A} \cdot \text{m}^{-2}$	$\text{C/s} \cdot \text{m}^2$	$\text{A/m}^2$

```
\dielectricconstant{\(magnitude)}
```

command	\dielectricconstant	
base	derived	alternate

N 2021-02-24

```
\direction{\langle magnitude \rangle}
\directionvector{\langle c_1, \dots, c_n \rangle}
\vector{direction}{\langle c_1, \dots, c_n \rangle}
```

command base	\direction derived	alternate
-----------------	-----------------------	-----------

N 2021-02-24

```
\displacement{\langle magnitude \rangle}
\displacementvector{\langle c_1, \dots, c_n \rangle}
\vector{displacement}{\langle c_1, \dots, c_n \rangle}
```

command base	\displacement derived	alternate
m	m	m

```
\duration{\langle magnitude \rangle}
```

command base	\duration derived	alternate
s	s	s

N 2021-02-24

```
\electricdipolemoment{\langle magnitude \rangle}
\electricdipolemomentvector{\langle c_1, \dots, c_n \rangle}
\vector{electricdipolemoment}{\langle c_1, \dots, c_n \rangle}
```

command base	\electricdipolemoment derived	alternate
A · s · m	C · m	C · m

N 2021-02-24

```
\electricfield{\langle magnitude \rangle}
\electricfieldvector{\langle c_1, \dots, c_n \rangle}
\vector{electricfield}{\langle c_1, \dots, c_n \rangle}
```

command base	\electricfield derived	alternate
kg · m · A <sup>-1</sup> · s <sup>-3</sup>	V/m	N/C

```
\electricflux{\langle magnitude \rangle}
```

command base	\electricflux derived	alternate
kg · m <sup>3</sup> · A <sup>-1</sup> · s <sup>-3</sup>	V · m	N · m <sup>2</sup> /C

```
\electricpotential{\langle magnitude \rangle}
```

<b>command</b>	\electricpotential	
<b>base</b>	derived	<b>alternate</b>
$\text{kg} \cdot \text{m}^2 \cdot \text{A}^{-1} \cdot \text{s}^{-3}$	V	V

N 2021-05-01

### \electricpotentialdifference{\(magnitude\)}

<b>command</b>	\electricpotentialdifference	
<b>base</b>	derived	<b>alternate</b>
$\text{kg} \cdot \text{m}^2 \cdot \text{A}^{-1} \cdot \text{s}^{-3}$	V	V

### \electroncurrent{\(magnitude\)}

<b>command</b>	\electroncurrent	
<b>base</b>	derived	<b>alternate</b>
$\text{s}^{-1}$	e/s	e/s

### \emf{\(magnitude\)}

<b>command</b>	\emf	
<b>base</b>	derived	<b>alternate</b>
$\text{kg} \cdot \text{m}^2 \cdot \text{A}^{-1} \cdot \text{s}^{-3}$	V	V

### \energy{\(magnitude\)}

<b>command</b>	\energy	
<b>base</b>	derived	<b>alternate</b>
$\text{kg} \cdot \text{m}^2 \cdot \text{s}^{-2}$	J	J

N 2021-04-15

### \energyinev{\(magnitude\)}

<b>command</b>	\energyinev	
<b>base</b>	derived	<b>alternate</b>
eV	eV	eV

N 2021-04-15

### \energyinkev{\(magnitude\)}

<b>command</b>	\energyinkev	
<b>base</b>	derived	<b>alternate</b>
keV	keV	keV

N 2021-04-15

### \energyinmev{\(magnitude\)}

<b>command</b>	\energyinmev	
<b>base</b>	derived	<b>alternate</b>
MeV	MeV	MeV

### \energydensity{\(magnitude\)}

<b>command</b>	<code>\energydensity</code>	
<b>base</b>	<code>derived</code>	<b>alternate</b>
$\text{kg} \cdot \text{m}^{-1} \cdot \text{s}^{-2}$	$\text{J/m}^3$	$\text{J/m}^3$

`\energyflux{<magnitude>}`  
`\energyfluxvector{<c1, ..., cn>}`  
`\vectorenergyflux{<c1, ..., cn>}`

<b>command</b>	<code>\energyflux</code>	
<b>base</b>	<code>derived</code>	<b>alternate</b>
$\text{kg} \cdot \text{s}^{-3}$	$\text{W/m}^2$	$\text{W/m}^2$

`\entropy{<magnitude>}`

<b>command</b>	<code>\entropy</code>	
<b>base</b>	<code>derived</code>	<b>alternate</b>
$\text{kg} \cdot \text{m}^2 \cdot \text{s}^{-2} \cdot \text{K}^{-1}$	$\text{J/K}$	$\text{J/K}$

`\force{<magnitude>}`  
`\forcevector{<c1, ..., cn>}`  
`\vectorforce{<c1, ..., cn>}`

<b>command</b>	<code>\force</code>	
<b>base</b>	<code>derived</code>	<b>alternate</b>
$\text{kg} \cdot \text{m} \cdot \text{s}^{-2}$	N	N

`\frequency{<magnitude>}`

<b>command</b>	<code>\frequency</code>	
<b>base</b>	<code>derived</code>	<b>alternate</b>
$\text{s}^{-1}$	Hz	Hz

`\gravitationalfield{<magnitude>}`  
`\gravitationalfieldvector{<c1, ..., cn>}`  
`\vectorgravitationalfield{<c1, ..., cn>}`

<b>command</b>	<code>\gravitationalfield</code>	
<b>base</b>	<code>derived</code>	<b>alternate</b>
$\text{m} \cdot \text{s}^{-2}$	N/kg	N/kg

`\gravitationalpotential{<magnitude>}`

<b>command</b>	<code>\gravitationalpotential</code>	
<b>base</b>	<code>derived</code>	<b>alternate</b>
$\text{m}^2 \cdot \text{s}^{-2}$	J/kg	J/kg

`\gravitationalpotentialdifference{<magnitude>}`

<b>command</b>	<code>\gravitationalpotentialdifference</code>	
<b>base</b>	<code>derived</code>	<b>alternate</b>
$\text{m}^2 \cdot \text{s}^{-2}$	$\text{J}/\text{kg}$	$\text{J}/\text{kg}$

`\impulse{\langle magnitude \rangle}`  
`\impulsevector{\langle c_1, \dots, c_n \rangle}`  
`\vectorimpulse{\langle c_1, \dots, c_n \rangle}`

<b>command</b>	<code>\impulse</code>	
<b>base</b>	<code>derived</code>	<b>alternate</b>
$\text{kg} \cdot \text{m} \cdot \text{s}^{-1}$	$\text{N} \cdot \text{s}$	$\text{N} \cdot \text{s}$

`\indexofrefraction{\langle magnitude \rangle}`

<b>command</b>	<code>\indexofrefraction</code>	
<b>base</b>	<code>derived</code>	<b>alternate</b>

`\inductance{\langle magnitude \rangle}`

<b>command</b>	<code>\inductance</code>	
<b>base</b>	<code>derived</code>	<b>alternate</b>
$\text{kg} \cdot \text{m}^2 \cdot \text{A}^{-2} \cdot \text{s}^{-2}$	$\text{H}$	$\text{V} \cdot \text{s}/\text{A}$

`\linearchargedensity{\langle magnitude \rangle}`

<b>command</b>	<code>\linearchargedensity</code>	
<b>base</b>	<code>derived</code>	<b>alternate</b>
$\text{A} \cdot \text{s} \cdot \text{m}^{-1}$	$\text{C}/\text{m}$	$\text{C}/\text{m}$

`\linearmassdensity{\langle magnitude \rangle}`

<b>command</b>	<code>\linearmassdensity</code>	
<b>base</b>	<code>derived</code>	<b>alternate</b>
$\text{kg} \cdot \text{m}^{-1}$	$\text{kg}/\text{m}$	$\text{kg}/\text{m}$

`\lorentzfactor{\langle magnitude \rangle}`

<b>command</b>	<code>\lorentzfactor</code>	
<b>base</b>	<code>derived</code>	<b>alternate</b>

`\luminousintensity{\langle magnitude \rangle}`

<b>command</b>	<code>\luminousintensity</code>	
<b>base</b>	<code>derived</code>	<b>alternate</b>
$\text{cd}$	$\text{cd}$	$\text{cd}$

N 2021-02-24

`\magneticcharge{magnitude}`

command	<code>\magneticcharge</code>	
base	<code>derived</code>	<code>alternate</code>
$A \cdot m$	$A \cdot m$	$A \cdot m$

`\magneticdipolemoment{magnitude}`

`\magneticdipolemomentvector{c1, ..., cn}`

`\vectormagneticdipolemoment{c1, ..., cn}`

command	<code>\magneticdipolemoment</code>	
base	<code>derived</code>	<code>alternate</code>
$A \cdot m^2$	$A \cdot m^2$	$J/T$

`\magneticfield{magnitude}`

`\magneticfieldvector{c1, ..., cn}`

`\vectormagneticfield{c1, ..., cn}`

command	<code>\magneticfield</code>	
base	<code>derived</code>	<code>alternate</code>
$kg \cdot A^{-1} \cdot s^{-2}$	$N/A \cdot m$	$T$

`\magneticflux{magnitude}`

command	<code>\magneticflux</code>	
base	<code>derived</code>	<code>alternate</code>
$kg \cdot m^2 \cdot A^{-1} \cdot s^{-2}$	$T \cdot m^2$	$V \cdot s$

`\mass{magnitude}`

command	<code>\mass</code>	
base	<code>derived</code>	<code>alternate</code>
$kg$	$kg$	$kg$

`\mobility{magnitude}`

command	<code>\mobility</code>	
base	<code>derived</code>	<code>alternate</code>
$kg \cdot m^2 \cdot A^{-1} \cdot s^{-4}$	$m^2/V \cdot s$	$C \cdot m/N \cdot s$

`\momentofinertia{magnitude}`

command	<code>\momentofinertia</code>	
base	<code>derived</code>	<code>alternate</code>
$kg \cdot m^2$	$J \cdot s^2$	$kg \cdot m^2$

N 2021-02-24

```
\momentum{\langle magnitude\rangle}
\momentumvector{\langle c_1, \dots, c_n \rangle}
\vectormomentum{\langle c_1, \dots, c_n \rangle}
```

<b>command</b>	<code>\momentum</code>	
<b>base</b>	<code>\derived</code>	<b>alternate</b>
$\text{kg} \cdot \text{m} \cdot \text{s}^{-1}$	$\text{kg} \cdot \text{m/s}$	$\text{kg} \cdot \text{m/s}$

N 2021-02-24

```
\momentumflux{\langle magnitude\rangle}
\momentumfluxvector{\langle c_1, \dots, c_n \rangle}
\vectormomentumflux{\langle c_1, \dots, c_n \rangle}
```

<b>command</b>	<code>\momentumflux</code>	
<b>base</b>	<code>\derived</code>	<b>alternate</b>
$\text{kg} \cdot \text{m}^{-1} \cdot \text{s}^{-2}$	$\text{N/m}^2$	$\text{N/m}^2$

```
\numberdensity{\langle magnitude\rangle}
```

<b>command</b>	<code>\numberdensity</code>	
<b>base</b>	<code>\derived</code>	<b>alternate</b>
$\text{m}^{-3}$	$/\text{m}^3$	$/\text{m}^3$

```
\permeability{\langle magnitude\rangle}
```

<b>command</b>	<code>\permeability</code>	
<b>base</b>	<code>\derived</code>	<b>alternate</b>
$\text{kg} \cdot \text{m} \cdot \text{A}^{-2} \cdot \text{s}^{-2}$	$\text{H/m}$	$\text{T} \cdot \text{m/A}$

```
\permittivity{\langle magnitude\rangle}
```

<b>command</b>	<code>\permittivity</code>	
<b>base</b>	<code>\derived</code>	<b>alternate</b>
$\text{A}^2 \cdot \text{s}^4 \cdot \text{kg}^{-1} \cdot \text{m}^{-3}$	$\text{F/m}$	$\text{C}^2/\text{N} \cdot \text{m}^2$

```
\planeangle{\langle magnitude\rangle}
```

<b>command</b>	<code>\planeangle</code>	
<b>base</b>	<code>\derived</code>	<b>alternate</b>
$\text{m} \cdot \text{m}^{-1}$	$\text{rad}$	$\text{rad}$

```
\polarizability{\langle magnitude\rangle}
```

<b>command</b>	<code>\polarizability</code>	
<b>base</b>	<code>\derived</code>	<b>alternate</b>
$\text{A}^2 \cdot \text{s}^4 \cdot \text{kg}^{-1}$	$\text{C} \cdot \text{m}^2/\text{V}$	$\text{C}^2 \cdot \text{m/N}$

```
\power{\langle magnitude\rangle}
```

<b>command</b>	<code>\power</code>	
<b>base</b>	<code>derived</code>	<b>alternate</b>
$\text{kg} \cdot \text{m}^2 \cdot \text{s}^{-3}$	<code>W</code>	$\text{J}/\text{s}$

`\poynting{<magnitude>}`  
`\poyntingvector{<c_1, ..., c_n>}`  
`\vectorpoynting{<c_1, ..., c_n>}`

<b>command</b>	<code>\poynting</code>	
<b>base</b>	<code>derived</code>	<b>alternate</b>
$\text{kg} \cdot \text{s}^{-3}$	<code>W/m<sup>2</sup></code>	$\text{W}/\text{m}^2$

`\pressure{<magnitude>}`

<b>command</b>	<code>\pressure</code>	
<b>base</b>	<code>derived</code>	<b>alternate</b>
$\text{kg} \cdot \text{m}^{-1} \cdot \text{s}^{-2}$	<code>Pa</code>	$\text{N}/\text{m}^2$

`\relativepermeability{<magnitude>}`

<b>command</b>	<code>\relativepermeability</code>	
<b>base</b>	<code>derived</code>	<b>alternate</b>

`\relativepermittivity{<magnitude>}`

<b>command</b>	<code>\relativepermittivity</code>	
<b>base</b>	<code>derived</code>	<b>alternate</b>

`\resistance{<magnitude>}`

<b>command</b>	<code>\resistance</code>	
<b>base</b>	<code>derived</code>	<b>alternate</b>
$\text{kg} \cdot \text{m}^2 \cdot \text{A}^{-2} \cdot \text{s}^{-3}$	$\Omega$	$\Omega$

`\resistivity{<magnitude>}`

<b>command</b>	<code>\resistivity</code>	
<b>base</b>	<code>derived</code>	<b>alternate</b>
$\text{kg} \cdot \text{m}^3 \cdot \text{A}^{-2} \cdot \text{s}^{-3}$	$\Omega \cdot \text{m}$	$\text{V} \cdot \text{m}/\text{A}$

`\solidangle{<magnitude>}`

<b>command</b>	<code>\solidangle</code>	
<b>base</b>	<code>derived</code>	<b>alternate</b>
$\text{m}^2 \cdot \text{m}^{-2}$	<code>sr</code>	$\text{sr}$

`\specificheatcapacity{magnitude}`

command	<code>\specificheatcapacity</code>	
base	<code>derived</code>	<code>alternate</code>
$\text{m}^2 \cdot \text{s}^{-2} \cdot \text{K}^{-1}$	$\text{J/K} \cdot \text{kg}$	$\text{J/K} \cdot \text{kg}$

`\springstiffness{magnitude}`

command	<code>\springstiffness</code>	
base	<code>derived</code>	<code>alternate</code>
$\text{kg} \cdot \text{s}^{-2}$	$\text{N/m}$	$\text{N/m}$

`\springstretch{magnitude}`

command	<code>\springstretch</code>	
base	<code>derived</code>	<code>alternate</code>
$\text{m}$	$\text{m}$	$\text{m}$

`\stress{magnitude}`

command	<code>\stress</code>	
base	<code>derived</code>	<code>alternate</code>
$\text{kg} \cdot \text{m}^{-1} \cdot \text{s}^{-2}$	$\text{Pa}$	$\text{N/m}^2$

`\strain{magnitude}`

command	<code>\strain</code>	
base	<code>derived</code>	<code>alternate</code>

`\temperature{magnitude}`

command	<code>\temperature</code>	
base	<code>derived</code>	<code>alternate</code>
$\text{K}$	$\text{K}$	$\text{K}$

`\torque{magnitude}`

`\torquevector{ $c_1, \dots, c_n$ }`

`\vectortorque{ $c_1, \dots, c_n$ }`

command	<code>\torque</code>	
base	<code>derived</code>	<code>alternate</code>
$\text{kg} \cdot \text{m}^2 \cdot \text{s}^{-2}$	$\text{N} \cdot \text{m}$	$\text{N} \cdot \text{m}$

`\velocity{magnitude}`

`\velocityvector{ $c_1, \dots, c_n$ }`

`\vectorvelocity{ $c_1, \dots, c_n$ }`

<b>command</b>	<code>\velocity</code>	
<b>base</b>	<code>derived</code>	<b>alternate</b>
$m \cdot s^{-1}$	$m/s$	$m/s$

`\velocityc{<magnitude>}`  
`\velocityvector{<c1, ..., cn>}`  
`\vectorvelocityc{<c1, ..., cn>}`

<b>command</b>	<code>\velocityc</code>	
<b>base</b>	<code>derived</code>	<b>alternate</b>
$c$	$c$	$c$

`\volume{<magnitude>}`

<b>command</b>	<code>\volume</code>	
<b>base</b>	<code>derived</code>	<b>alternate</b>
$m^3$	$m^3$	$m^3$

`\volumechargedensity{<magnitude>}`

<b>command</b>	<code>\volumechargedensity</code>	
<b>base</b>	<code>derived</code>	<b>alternate</b>
$A \cdot s/m^3$	$C/m^3$	$C/m^3$

`\volumemassdensity{<magnitude>}`

<b>command</b>	<code>\volumemassdensity</code>	
<b>base</b>	<code>derived</code>	<b>alternate</b>
$kg \cdot m^{-3}$	$kg/m^3$	$kg/m^3$

`\wavelength{<magnitude>}`

<b>command</b>	<code>\wavelength</code>	
<b>base</b>	<code>derived</code>	<b>alternate</b>
$m$	$m$	$m$

`\wavenumber{<magnitude>}`  
`\wavenumbervector{<c1, ..., cn>}`  
`\vectorwavenumber{<c1, ..., cn>}`

<b>command</b>	<code>\wavenumber</code>	
<b>base</b>	<code>derived</code>	<b>alternate</b>
$m^{-1}$	$/m$	$/m$

`\work{<magnitude>}`

<b>command</b>	<code>\work</code>	
<b>base</b>	<code>derived</code>	<b>alternate</b>
$\text{kg} \cdot \text{m}^2 \cdot \text{s}^{-2}$	J	J

```
\youngsmodulus{(magnitude)}
```

<b>command</b>	<code>\youngsmodulus</code>	
<b>base</b>	<code>derived</code>	<b>alternate</b>
$\text{kg} \cdot \text{m}^{-1} \cdot \text{s}^{-2}$	Pa	$\text{N}/\text{m}^2$

#### 4.4.4 Defining and Redefining Physical Quantities

N 2021-02-16  
N 2021-02-21

```
\newscalarquantity{(name)}{(base units)}[(derived units)][(alternate units)]
\renewscalearquantity{(name)}{(base units)}[(derived units)][(alternate units)]
```

Command to (re)define a new/existing scalar quantity. If the derived or alternate units are omitted, they are defined to be the same as the base units. Do not use both this command and `\newvectorquantity` or `\renewvectorquantity` to (re)define a quantity.

N 2021-02-16  
N 2021-02-21

```
\newvectorquantity{(name)}{(base units)}[(derived units)][(alternate units)]
\renewvectorquantity{(name)}{(base units)}[(derived units)][(alternate units)]
```

Command to (re)define a new/existing vector quantity. If the derived or alternate units are omitted, they are defined to be the same as the base units. Do not use both this command and `\newscalarquantity` or `\renewscalearquantity` to (re)define a quantity.

#### 4.4.5 Changing Units

Units are set when `mandi` is loaded, but the default setting can be easily overridden in four ways: command variants that are defined when a [physical quantity](#) <sup>P.9</sup> or [physical constant](#) <sup>P.23</sup> is defined, a global modal command (switch), a command that sets units for a single instance, and an environment that sets units for its duration. All of these methods work for both physical quantities and physical constants.

U 2021-02-26  
U 2021-02-26  
U 2021-02-26

```
\alwaysusebaseunits
\alwaysusederivedunits
\alwaysusealternateunits
```

Modal commands (switches) for setting the default unit form for the entire document. When `mandi` is loaded, one of these three commands is executed depending on whether the optional `units` key is provided. See the section on loading the package for details. Alternate units are the default because they are the most likely ones to be seen in introductory physics textbooks.

U 2021-02-26  
U 2021-02-26  
U 2021-02-26

```
\hereusebaseunits{(content)}
\hereusederivedunits{(content)}
\hereusedalternateunits{(content)}
```

Commands for setting the unit form on the fly for a single instance. The example uses momentum and the Coulomb constant, but they work for any defined quantity and constant.

\(\hereusebaseunits{\momentum{5}}\)	\(\hereusederivedunits{\momentum{5}}\)	\(\hereusealternateunits{\momentum{5}}\)	\(\hereusebaseunits{\oofpez}\)	\(\hereusederivedunits{\oofpez}\)	\(\hereusealternateunits{\oofpez}\)
\(\hereusebaseunits{\momentum{5}}\)	\(\hereusederivedunits{\momentum{5}}\)	\(\hereusealternateunits{\momentum{5}}\)	\(\hereusebaseunits{\oofpez}\)	\(\hereusederivedunits{\oofpez}\)	\(\hereusealternateunits{\oofpez}\)
\(\hereusebaseunits{\momentum{5}}\)	\(\hereusederivedunits{\momentum{5}}\)	\(\hereusealternateunits{\momentum{5}}\)	\(\hereusebaseunits{\oofpez}\)	\(\hereusederivedunits{\oofpez}\)	\(\hereusealternateunits{\oofpez}\)
\(\hereusebaseunits{\momentum{5}}\)	\(\hereusederivedunits{\momentum{5}}\)	\(\hereusealternateunits{\momentum{5}}\)	\(\hereusebaseunits{\oofpez}\)	\(\hereusederivedunits{\oofpez}\)	\(\hereusealternateunits{\oofpez}\)
\(\hereusebaseunits{\momentum{5}}\)	\(\hereusederivedunits{\momentum{5}}\)	\(\hereusealternateunits{\momentum{5}}\)	\(\hereusebaseunits{\oofpez}\)	\(\hereusederivedunits{\oofpez}\)	\(\hereusealternateunits{\oofpez}\)

U 2021-02-26

```
\begin{usebaseunits}
  (environment content)
\end{usebaseunits}
```

(use base units)

U 2021-02-26

```
\begin{usederivedunits}
  (environment content)
\end{usederivedunits}
```

(use derived units)

U 2021-02-26

```
\begin{usealternateunits}
  (environment content)
\end{usealternateunits}
```

(use alternate units)

Inside these environments units are changed for the duration of the environment regardless of the global default setting.

\(\momentum{5}\)	\(\oofpez\)	\(\usebaseunits{\momentum{5}}\)	\(\usebaseunits{\oofpez}\)	\(\usederivedunits{\momentum{5}}\)	\(\usederivedunits{\oofpez}\)	\(\usealternateunits{\momentum{5}}\)	\(\usealternateunits{\oofpez}\)
\(\momentum{5}\)	\(\oofpez\)	\(\usebaseunits{\momentum{5}}\)	\(\usebaseunits{\oofpez}\)	\(\usederivedunits{\momentum{5}}\)	\(\usederivedunits{\oofpez}\)	\(\usealternateunits{\momentum{5}}\)	\(\usealternateunits{\oofpez}\)
\(\momentum{5}\)	\(\oofpez\)	\(\usebaseunits{\momentum{5}}\)	\(\usebaseunits{\oofpez}\)	\(\usederivedunits{\momentum{5}}\)	\(\usederivedunits{\oofpez}\)	\(\usealternateunits{\momentum{5}}\)	\(\usealternateunits{\oofpez}\)
\(\momentum{5}\)	\(\oofpez\)	\(\usebaseunits{\momentum{5}}\)	\(\usebaseunits{\oofpez}\)	\(\usederivedunits{\momentum{5}}\)	\(\usederivedunits{\oofpez}\)	\(\usealternateunits{\momentum{5}}\)	\(\usealternateunits{\oofpez}\)
\(\momentum{5}\)	\(\oofpez\)	\(\usebaseunits{\momentum{5}}\)	\(\usebaseunits{\oofpez}\)	\(\usederivedunits{\momentum{5}}\)	\(\usederivedunits{\oofpez}\)	\(\usealternateunits{\momentum{5}}\)	\(\usealternateunits{\oofpez}\)
\(\momentum{5}\)	\(\oofpez\)	\(\usebaseunits{\momentum{5}}\)	\(\usebaseunits{\oofpez}\)	\(\usederivedunits{\momentum{5}}\)	\(\usederivedunits{\oofpez}\)	\(\usealternateunits{\momentum{5}}\)	\(\usealternateunits{\oofpez}\)

## 4.5 Physical Constants

### 4.5.1 Typesetting Physical Constants

Take the quantity  $\frac{1}{4\pi\epsilon_0}$ , sometimes called the [Coulomb constant](#), as the prototypical [physical constant](#) in an introductory physics course. Here are all the ways to access this quantity in `mandi`. As you can see, these commands are almost identical to the corresponding commands for physical quantities.

`\oofpez`

Command for the Coulomb constant. The constant's numerical precision and default units will depend on the options passed to `mandi` at load time. Alternate units and approximate numerical values are the defaults. Other units can be forced as demonstrated.

<code>\(\backslash\oofpez\)</code>	<code>\backslash\oofpez</code>	$9 \times 10^9 \text{ N} \cdot \text{m}^2/\text{C}^2$
<code>\(\backslash\oofpezapproximatevalue\)</code>	<code>\backslash\oofpezapproximatevalue</code>	$9 \times 10^9$
<code>\(\backslash\oofpezprecisevalue\)</code>	<code>\backslash\oofpezprecisevalue</code>	$8.9875517923 \times 10^9$
<code>\(\backslash\oofpezmathsymbol\)</code>	<code>\backslash\oofpezmathsymbol</code>	$\frac{1}{4\pi\epsilon_0}$
<code>\(\backslash\oofpezbaseunits\)</code>	<code>\backslash\oofpezbaseunits</code>	$9 \times 10^9 \text{ kg} \cdot \text{m}^3 \cdot \text{A}^{-2} \cdot \text{s}^{-4}$
<code>\(\backslash\oofpezderivedunits\)</code>	<code>\backslash\oofpezderivedunits</code>	$9 \times 10^9 \text{ m/F}$
<code>\(\backslash\oofpezelternateunits\)</code>	<code>\backslash\oofpezelternateunits</code>	$9 \times 10^9 \text{ N} \cdot \text{m}^2/\text{C}^2$
<code>\(\backslash\oofpezonlybaseunits\)</code>	<code>\backslash\oofpezonlybaseunits</code>	$\text{kg} \cdot \text{m}^3 \cdot \text{A}^{-2} \cdot \text{s}^{-4}$
<code>\(\backslash\oofpezonlyderivedunits\)</code>	<code>\backslash\oofpezonlyderivedunits</code>	$\text{m/F}$
<code>\(\backslash\oofpezonlyalternateunits\)</code>	<code>\backslash\oofpezonlyalternateunits</code>	$\text{N} \cdot \text{m}^2/\text{C}^2$

## 4.5.2 Checking Physical Constants

U 2022-01-27

### `\checkconstant{\(name\)}`

Command to check and typeset the constant's name, mathematical symbol, approximate value, precise value, base units, derived units, and alternate units.

## 4.5.3 Predefined Physical Constants

Every other defined physical constant can be treated similarly to `\oofpez`<sup>P.27</sup>. Just replace `\oofpez` with the constant's name. Unfortunately, there is no universal agreement on the names of every constant so don't fret if the names used here vary from other sources. Here are all the physical constants, with all their units, defined in `mandi`. The constants `\coulombconstant`<sup>P.25</sup> and `\biotsavartconstant` are defined as semantic aliases for, respectively, `\oofpez`<sup>P.27</sup> and `\mzofp`<sup>P.27</sup>.

### `\avogadro`

(exact)

<b>command</b>	<code>\avogadro</code>	
<b>symbol</b>	<code>approximate</code>	<code>precise</code>
$N_A$	$6 \times 10^{23}$	$6.02214076 \times 10^{23}$
<b>base</b>	<code>derived</code>	<code>alternate</code>
$\text{mol}^{-1}$	<code>/mol</code>	<code>/mol</code>

N 2021-02-02

### `\biotsavartconstant`

<b>command</b>	<code>\biotsavartconstant</code>	
<b>symbol</b>	<code>approximate</code>	<code>precise</code>
$\frac{\mu_0}{4\pi}$	$10^{-7}$	$10^{-7}$
<b>base</b>	<code>derived</code>	<code>alternate</code>
$\text{kg} \cdot \text{m} \cdot \text{A}^{-2} \cdot \text{s}^{-2}$	<code>H/m</code>	$\text{T} \cdot \text{m/A}$

### `\bohrradius`

**command**  
**symbol**  
 $a_0$   
**base**  
m

\bohradius  
**approximate**  
 $5.3 \times 10^{-11}$   
**derived**  
m

**precise**  
 $5.29177210903 \times 10^{-11}$   
**alternate**  
m

### \boltzmann

(exact)

**command**  
**symbol**  
 $k_B$   
**base**  
 $kg \cdot m^2 \cdot s^{-2} \cdot K^{-1}$

\boltzmann  
**approximate**  
 $1.4 \times 10^{-23}$   
**derived**  
J/K

**precise**  
 $1.380649 \times 10^{-23}$   
**alternate**  
J/K

N 2021-02-02

### \coulombconstant

**command**  
**symbol**  
 $\frac{1}{4\pi\epsilon_0}$   
**base**  
 $kg \cdot m^3 \cdot A^{-2} \cdot s^{-4}$

\coulombconstant  
**approximate**  
 $9 \times 10^9$   
**derived**  
m/F

**precise**  
 $8.9875517923 \times 10^9$   
**alternate**  
 $N \cdot m^2/C^2$

### \earthmass

**command**  
**symbol**  
 $M_{Earth}$   
**base**  
kg

\earthmass  
**approximate**  
 $6.0 \times 10^{24}$   
**derived**  
kg

**precise**  
 $5.9722 \times 10^{24}$   
**alternate**  
kg

### \earthmoondistance

**command**  
**symbol**  
 $d_{EM}$   
**base**  
m

\earthmoondistance  
**approximate**  
 $3.8 \times 10^8$   
**derived**  
m

**precise**  
 $3.81550 \times 10^8$   
**alternate**  
m

### \earthradius

**command**  
**symbol**  
 $R_{Earth}$   
**base**  
m

\earthradius  
**approximate**  
 $6.4 \times 10^6$   
**derived**  
m

**precise**  
 $6.3781 \times 10^6$   
**alternate**  
m

### \earthsundistance

<b>command</b>	\earthsundistance	
<b>symbol</b>	approximate	<b>precise</b>
$d_{ES}$	$1.5 \times 10^{11}$	$1.496 \times 10^{11}$
<b>base</b>	derived	alternate
m	m	m

### \electroncharge

<b>command</b>	\electroncharge	
<b>symbol</b>	approximate	<b>precise</b>
$q_e$	$-1.6 \times 10^{-19}$	$-1.602176634 \times 10^{-19}$
<b>base</b>	derived	alternate
$A \cdot s$	C	C

### \electronCharge

<b>command</b>	\electronCharge	
<b>symbol</b>	approximate	<b>precise</b>
$Q_e$	$-1.6 \times 10^{-19}$	$-1.602176634 \times 10^{-19}$
<b>base</b>	derived	alternate
$A \cdot s$	C	C

### \electronmass

<b>command</b>	\electronmass	
<b>symbol</b>	approximate	<b>precise</b>
$m_e$	$9.1 \times 10^{-31}$	$9.1093837015 \times 10^{-31}$
<b>base</b>	derived	alternate
kg	kg	kg

### \elementarycharge

(exact)

<b>command</b>	\elementarycharge	
<b>symbol</b>	approximate	<b>precise</b>
e	$1.6 \times 10^{-19}$	$1.602176634 \times 10^{-19}$
<b>base</b>	derived	alternate
$A \cdot s$	C	C

### \finestructure

<b>command</b>	\finestructure	
<b>symbol</b>	approximate	<b>precise</b>
$\alpha$	$\frac{1}{137}$	$7.2973525693 \times 10^{-3}$
<b>base</b>	derived	alternate

### \hydrogenmass

<b>command</b>	\hydrogenmass	
<b>symbol</b>	approximate	<b>precise</b>
$m_H$	$1.7 \times 10^{-27}$	$1.6737236 \times 10^{-27}$
<b>base</b>	derived	alternate
kg	kg	kg

### \moonearthdistance

<b>command</b>	\moonearthdistance	
<b>symbol</b>	approximate	<b>precise</b>
$d_{ME}$	$3.8 \times 10^8$	$3.81550 \times 10^8$
<b>base</b>	derived	alternate
m	m	m

### \moonmass

<b>command</b>	\moonmass	
<b>symbol</b>	approximate	<b>precise</b>
$M_{Moon}$	$7.3 \times 10^{22}$	$7.342 \times 10^{22}$
<b>base</b>	derived	alternate
kg	kg	kg

### \moonradius

<b>command</b>	\moonradius	
<b>symbol</b>	approximate	<b>precise</b>
$R_{Moon}$	$1.7 \times 10^6$	$1.7371 \times 10^6$
<b>base</b>	derived	alternate
m	m	m

### \mzofp

<b>command</b>	\mzofp	
<b>symbol</b>	approximate	<b>precise</b>
$\frac{\mu_0}{4\pi}$	$10^{-7}$	$10^{-7}$
<b>base</b>	derived	alternate
$\text{kg} \cdot \text{m} \cdot \text{A}^{-2} \cdot \text{s}^{-2}$	H/m	$\text{T} \cdot \text{m/A}$

### \neutronmass

<b>command</b>	\neutronmass	
<b>symbol</b>	approximate	<b>precise</b>
$m_n$	$1.7 \times 10^{-27}$	$1.67492749804 \times 10^{-27}$
<b>base</b>	derived	alternate
kg	kg	kg

### \oofpez

<b>command</b>	\oofpez	
<b>symbol</b>	approximate	<b>precise</b>
$\frac{1}{4\pi\epsilon_0}$	$9 \times 10^9$	$8.9875517923 \times 10^9$
<b>base</b>	derived	<b>alternate</b>
$\text{kg} \cdot \text{m}^3 \cdot \text{A}^{-2} \cdot \text{s}^{-4}$	m/F	$\text{N} \cdot \text{m}^2/\text{C}^2$

### \oofpezcs

<b>command</b>	\oofpezcs	
<b>symbol</b>	approximate	<b>precise</b>
$\frac{1}{4\pi\epsilon_0 c^2}$	$10^{-7}$	$10^{-7}$
<b>base</b>	derived	<b>alternate</b>
$\text{kg} \cdot \text{m} \cdot \text{A}^{-2} \cdot \text{s}^{-2}$	$\text{T} \cdot \text{m}^2$	$\text{N} \cdot \text{s}^2/\text{C}^2$

### \planck

(exact)

<b>command</b>	\planck	
<b>symbol</b>	approximate	<b>precise</b>
h	$6.6 \times 10^{-34}$	$6.62607015 \times 10^{-34}$
<b>base</b>	derived	<b>alternate</b>
$\text{kg} \cdot \text{m}^2 \cdot \text{s}^{-1}$	J · s	J · s

### \planckbar

<b>command</b>	\planckbar	
<b>symbol</b>	approximate	<b>precise</b>
$\hbar$	$1.1 \times 10^{-34}$	$1.054571817 \times 10^{-34}$
<b>base</b>	derived	<b>alternate</b>
$\text{kg} \cdot \text{m}^2 \cdot \text{s}^{-1}$	J · s	J · s

### \planckc

<b>command</b>	\planckc	
<b>symbol</b>	approximate	<b>precise</b>
hc	$2.0 \times 10^{-25}$	$1.98644586 \times 10^{-25}$
<b>base</b>	derived	<b>alternate</b>
$\text{kg} \cdot \text{m}^3 \cdot \text{s}^{-2}$	J · m	J · m

### \protoncharge

<b>command</b>	\protoncharge	
<b>symbol</b>	approximate	<b>precise</b>
$q_p$	$+1.6 \times 10^{-19}$	$+1.602176634 \times 10^{-19}$
<b>base</b>	derived	<b>alternate</b>
$\text{A} \cdot \text{s}$	C	C

### \protonCharge

**command**  
**symbol**  
 $Q_p$   
**base**  
 $A \cdot s$

\protonCharge  
**approximate**  
 $+1.6 \times 10^{-19}$   
**derived**  
C

**precise**  
 $+1.602176634 \times 10^{-19}$   
**alternate**  
C

### \protonmass

**command**  
**symbol**  
 $m_p$   
**base**  
kg

\protonmass  
**approximate**  
 $1.7 \times 10^{-27}$   
**derived**  
kg

**precise**  
 $1.672621898 \times 10^{-27}$   
**alternate**  
kg

### \rydberg

**command**  
**symbol**  
 $R_\infty$   
**base**  
 $m^{-1}$

\rydberg  
**approximate**  
 $1.1 \times 10^7$   
**derived**  
 $m^{-1}$

**precise**  
 $1.0973731568160 \times 10^7$   
**alternate**  
 $m^{-1}$

### \speedoflight

(exact)

**command**  
**symbol**  
c  
**base**  
 $m \cdot s^{-1}$

\speedoflight  
**approximate**  
 $3 \times 10^8$   
**derived**  
m/s

**precise**  
 $2.99792458 \times 10^8$   
**alternate**  
m/s

### \stefanboltzmann

**command**  
**symbol**  
 $\sigma$   
**base**  
 $kg \cdot s^{-3} \cdot K^{-4}$

\stefanboltzmann  
**approximate**  
 $5.7 \times 10^{-8}$   
**derived**  
 $W/m^2 \cdot K^4$

**precise**  
 $5.670374 \times 10^{-8}$   
**alternate**  
 $W/m^2 \cdot K^4$

### \sunearthdistance

**command**  
**symbol**  
 $d_{SE}$   
**base**  
m

\sunearthdistance  
**approximate**  
 $1.5 \times 10^{11}$   
**derived**  
m

**precise**  
 $1.496 \times 10^{11}$   
**alternate**  
m

### \sunradius

<b>command</b>	\sunradius	
<b>symbol</b>	approximate	<b>precise</b>
$R_{\text{Sun}}$	$7.0 \times 10^8$	$6.957 \times 10^8$
<b>base</b>	derived	alternate
m	m	m

### \surfacegravfield

<b>command</b>	\surfacegravfield	
<b>symbol</b>	approximate	<b>precise</b>
g	9.8	9.807
<b>base</b>	derived	alternate
$\text{m} \cdot \text{s}^{-2}$	N/kg	N/kg

### \universalgrav

<b>command</b>	\universalgrav	
<b>symbol</b>	approximate	<b>precise</b>
G	$6.7 \times 10^{-11}$	$6.67430 \times 10^{-11}$
<b>base</b>	derived	alternate
$\text{m}^3 \cdot \text{kg}^{-1} \cdot \text{s}^{-2}$	$\text{N} \cdot \text{m}^2/\text{kg}^2$	$\text{N} \cdot \text{m}^2/\text{kg}^2$

### \vacuumpermeability

<b>command</b>	\vacuumpermeability	
<b>symbol</b>	approximate	<b>precise</b>
$\mu_0$	$4\pi \times 10^{-7}$	$4\pi \times 10^{-7}$
<b>base</b>	derived	alternate
$\text{kg} \cdot \text{m} \cdot \text{A}^{-2} \cdot \text{s}^{-2}$	H/m	T · m/A

### \vacuumpermittivity

<b>command</b>	\vacuumpermittivity	
<b>symbol</b>	approximate	<b>precise</b>
$\epsilon_0$	$9 \times 10^{-12}$	$8.854187817 \times 10^{-12}$
<b>base</b>	derived	alternate
$\text{A}^2 \cdot \text{s}^4 \cdot \text{kg}^{-1} \cdot \text{m}^{-3}$	F/m	$\text{C}^2/\text{N} \cdot \text{m}^2$

#### 4.5.4 Defining and Redefining Physical Constants

N 2021-02-16 `\newphysicalconstant {<name>} {<symbol>} {<approximate value>} {<precise value>} {<base units>} [<derived units>] [<alternate units>]`  
N 2021-02-21 `\renewphysicalconstant {<name>} {<symbol>} {<approximate value>} {<precise value>} {<base units>} [<derived units>] [<alternate units>]`

Command to define/redefine a new/existing physical constant. If the derived or alternate units are omitted, they are defined to be the same as the base units.

#### 4.5.5 Changing Precision

[Changing units](#) <sup>P.22</sup> works for physical constants just as it does for physical quantities. A similar mechanism is provided for changing the precision of physical constants' numerical values.

N 2021-02-16  
N 2021-02-16

`\alwaysuseapproximateconstants`  
`\alwaysusepreciseconstants`

Modal commands (switches) for setting the default precision for the entire document. The default when the package is loaded is set by the presence or absence of the [preciseconstants](#) <sup>P.8</sup> key.

N 2021-02-16  
N 2021-02-16

`\hereuseapproximateconstants{<content>}`  
`\hereusepreciseconstants{<content>}`

Commands for setting the precision on the fly for a single instance.

```
\(\hereuseapproximateconstants{\oofpez} \) \\
```

$9 \times 10^9 \text{ N} \cdot \text{m}^2/\text{C}^2$   
 $8.9875517923 \times 10^9 \text{ N} \cdot \text{m}^2/\text{C}^2$

N 2021-02-16  
N 2021-02-16

`\begin{useapproximateconstants}`  
  *(environment content)*  
`\end{useapproximateconstants}`  
`\begin{usepreciseconstants}`  
  *(environment content)*  
`\end{usepreciseconstants}`

(use approximate constants)

(use precise constants)

Inside these environments precision is changed for the duration of the environment regardless of the global default setting.

```
\(\oofpez \) \\
```

```
\begin{useapproximateconstants}
```

```
\(\oofpez \) \\
```

```
\end{useapproximateconstants}
```

```
\begin{usepreciseconstants}
```

```
\(\oofpez \) \\
```

```
\end{usepreciseconstants}
```

```
\(\oofpez \)
```

$9 \times 10^9 \text{ N} \cdot \text{m}^2/\text{C}^2$   
 $9 \times 10^9 \text{ N} \cdot \text{m}^2/\text{C}^2$   
 $8.9875517923 \times 10^9 \text{ N} \cdot \text{m}^2/\text{C}^2$   
 $9 \times 10^9 \text{ N} \cdot \text{m}^2/\text{C}^2$

#### 4.6 Predefined Units and Constructs

These commands should be used only in defining or redefining physical quantities or physical constants. One exception is `\emptyunit`, which may be used for explanatory purposes.

`\per`  
`\usk`  
`\unit{<magnitude>}{<unit>}`  
`\emptyunit`  
`\ampere`

<u>N</u> 2021-04-15	\atomicmassunit	
	\candela	(not SI but common in introductory physics)
	\coulomb	(alias)
	\degree	
	\electronvolt	
	\ev	
	\farad	
	\henry	
	\hertz	
	\joule	
	\kelvin	
	\kev	(alias)
<u>N</u> 2021-04-15	\kilolectronvolt	(not SI but common in introductory physics)
<u>N</u> 2021-04-15	\kilogram	
	\lightspeed	(not SI but common relativity)
	\megaelectronvolt	(not SI but common in introductory physics)
	\meter	
	\metre	
	\mev	(alias)
<u>N</u> 2021-04-15	\mole	(alias)
	\newton	
	\ohm	
	\pascal	
	\radian	
	\second	
	\siemens	
	\steradian	
	\tesla	
	\volt	
	\watt	
	\weber	
	\tothetwo	(postfix)
	\tothethree	(postfix)
	\tothefour	(postfix)
	\inverse	(postfix)
	\totheinversetwo	(postfix)
	\totheinversethree	(postfix)
	\totheinversefour	(postfix)

	/
	.
	3 m/s
\(\per\)	\( \per \)
\(\usk\)	\( \usk \)
\(\unit{3}{\meter\per\second}\)	\( \unit{3}{\meter\per\second} \)
\(\emptyunit\)	\( \emptyunit \)
\(\ampere\)	\( \ampere \)
\(\atomicmassunit\)	\( \atomicmassunit \)
\(\candela\)	\( \candela \)
\(\coulomb\)	\( \coulomb \)
\(\degree\)	\( \degree \)
\(\electronvolt\)	\( \electronvolt \)
\(\farad\)	\( \farad \)
\(\henry\)	\( \henry \)
\(\hertz\)	\( \hertz \)
\(\joule\)	\( \joule \)
\(\kelvin\)	\( \kelvin \)
\(\kev\)	\( \kev \)
\(\kilogram\)	\( \kilogram \)
\(\lightspeed\)	\( \lightspeed \)
\(\meter\)	\( \meter \)
\(\metre\)	\( \metre \)
\(\mev\)	\( \mev \)
\(\mole\)	\( \mole \)
\(\newton\)	\( \newton \)
\(\ohm\)	\( \ohm \)
\(\pascal\)	\( \pascal \)
\(\radian\)	\( \radian \)
\(\second\)	\( \second \)
\(\siemens\)	\( \siemens \)
\(\steradian\)	\( \steradian \)
\(\tesla\)	\( \tesla \)
\(\volt\)	\( \volt \)
\(\watt\)	\( \watt \)
\(\weber\)	\( \weber \)
\(\emptyunit{2}\)	\( \emptyunit{2} \)
\(\emptyunit{3}\)	\( \emptyunit{3} \)
\(\emptyunit{4}\)	\( \emptyunit{4} \)
\(\emptyunit{\inverse}\)	\( \emptyunit{\inverse} \)
\(\emptyunit{\inversetwo}\)	\( \emptyunit{\inversetwo} \)
\(\emptyunit{\inversethree}\)	\( \emptyunit{\inversethree} \)
\(\emptyunit{\inversefour}\)	\( \emptyunit{\inversefour} \)
	\( \square \)
	A
	u
	cd
	C
	°
	eV
	F
	H
	Hz
	J
	K
	keV
	kg
	c
	m
	m
	MeV
	mol
	N
	Ω
	Pa
	rad
	s
	S
	sr
	T
	V
	W
	Wb
	□²
	□³
	□⁴
	□⁻¹
	□⁻²
	□⁻³
	□⁻⁴

2022-01-27

## \hbar

A better glyph for Planck's constant over  $2\pi$ .

\( \backslash \hbar \)

1

```
\tent{<number>}\timestento{<number>}\xtento{<number>}
```

Commands for powers of ten and scientific notation.

```
\( \tent{4} \) \\  
\( 3\tent{8} \) \\  
\( 3\xtent{8} \)
```

$$10^{-4}$$

U 2022-01-27

**\mivector** [ $\langle \text{delimiter} \rangle$ ] { $\langle c_1, \dots, c_n \rangle$ } [ $\langle \text{units} \rangle$ ]

Typesets a vector as either numeric or symbolic components with an optional unit (for numerical components only). There must be more than one component, and there can be more than three components. The delimiter used in the list of components can be specified; the default is a comma. The notation mirrors that of *Matter & Interactions*.

```

\\( \mivector{p_0,p_1,p_2,p_3} \\) //  $\langle p_0, p_1, p_2, p_3 \rangle$ 
\\( \mivector{\gamma mc,\gamma mv_x,\gamma mv_y,\gamma mv_z} \\) //  $\langle \gamma mc, \gamma mv_x, \gamma mv_y, \gamma mv_z \rangle$ 
\\( \mivector{\frac{Q_1 Q_2}{x^2},0,0} \\) //  $\left\langle \frac{Q_1 Q_2}{x^2}, 0, 0 \right\rangle$ 
\\( \mivector{-1,0,0} \\) //  $\langle -1, 0, 0 \rangle$ 
\\( \mivector{-1,0,0}[\velocityonlyderivedunits] \\) //  $\langle -1, 0, 0 \rangle$  m/s
\\( \mivector{-1,0,0}[\meter\per\second] \\) //  $\langle -1, 0, 0 \rangle$  m/s
\\( \velocity{\mivector{-1,0,0}} \\) //  $\langle -1, 0, 0 \rangle$  m/s

```

## 4.7 **mandi** Source Code

Define the package version and date for global use, exploiting the fact that in a .sty file there is now no need for `\makeatletter` and `\makeatother`. This simplifies defining internal commands (with @ in the name) that are not for the user to know about.

```

1 \def\mandi@version{3.1.0}
2 \def\mandi@date{2022-01-27}
3 \NeedsTeXFormat{LaTeXe}[2020-02-02]
4 \DeclareRelease{v3.1.0}{2022-01-27}{mandi.sty}
5 \DeclareCurrentRelease{v\mandi@version}{\mandi@date}
6 \ProvidesPackage{mandi}
7   [\mandi@date\space v\mandi@version\space Macros for physical quantities]
```

Define a convenient package version command.

```
8 \newcommand*\mandiversion{v\mandi@version\space dated \mandi@date}
```

Load third party packages, documenting why each one is needed.

```

9 \RequirePackage{pgfopts}          % needed for key-value interface
10 \RequirePackage{array}           % needed for \checkquantity and \checkconstant
11 \RequirePackage{iftex}            % needed for requiring LuaLaTeX
12 \RequirePackage{unicode-math}     % needed for Unicode support

13 \IfFormatAtLeastTF {2020-10-01} % load xparse if necessary
14 {}%
15 {\RequirePackage{xparse}}%
16 \RequireLuaTeX                  % require this engine
```

Parts of the unit engine have been rewritten with `xparse` for both clarity and power. Note that `xparse` is now part of the L<sup>A</sup>T<sub>E</sub>X 2 <sub>$\epsilon$</sub>  kernel. Other parts have been rewritten in `expl` with a look to the future.

Generic internal selectors.

```
17 \newcommand*\mandi@selectunits(){}
18 \newcommand*\mandi@selectprecision{}}
```

Specific internal selectors.

```

19 \newcommand*\mandi@selectapproximate[2]{#1}      % really \@firstoftwo
20 \newcommand*\mandi@selectprecise[2]{#2}           % really \@secondoftwo
21 \newcommand*\mandi@selectbaseunits[3]{#1}          % really \@firstofthree
22 \newcommand*\mandi@selectderivedunits[3]{#2}        % really \@secondofthree
23 \newcommand*\mandi@selectalternateunits[3]{#3}      % really \@thirdofthree
```

Document level global switches.

```

24 \NewDocumentCommand{\alwaysusebaseunits} {}
25   {\renewcommand*\mandi@selectunits{\mandi@selectbaseunits}}%
26 \NewDocumentCommand{\alwaysusederivedunits} {}
27   {\renewcommand*\mandi@selectunits{\mandi@selectderivedunits}}%
28 \NewDocumentCommand{\alwaysusealternateunits} {}
29   {\renewcommand*\mandi@selectunits{\mandi@selectalternateunits}}%
30 \NewDocumentCommand{\alwaysuseapproximateconstants} {}
31   {\renewcommand*\mandi@selectprecision{\mandi@selectapproximate}}%
32 \NewDocumentCommand{\alwaysusepreciseconstants} {}
33   {\renewcommand*\mandi@selectprecision{\mandi@selectprecise}}%
```

Document level localized variants.

```

34 \NewDocumentCommand{\hereusebaseunits}{ m }{\begingroup\alwaysusebaseunits#1\endgroup}%
35 \NewDocumentCommand{\hereusederivedunits}{ m }{\begingroup\alwaysusederivedunits#1\endgroup}%
36 \NewDocumentCommand{\hereusealternateunits}{ m }{\begingroup\alwaysusealternateunits#1\endgroup}%
37 \NewDocumentCommand{\hereuseapproximateconstants}{ m }{\begingroup\alwaysuseapproximateconstants#1\endgroup}%
38 \NewDocumentCommand{\hereusepreciseconstants}{ m }{\begingroup\alwaysusepreciseconstants#1\endgroup}%

```

Document level environments.

```
39 \NewDocumentEnvironment{usebaseunits}{}{\alwaysusebaseunits}%
40 \NewDocumentEnvironment{usederivedunits}{}{\alwaysusederivedunits}%
41 \NewDocumentEnvironment{usealternateunits}{}{\alwaysusealternateunits}%
42 \NewDocumentEnvironment{useapproximateconstants}{}{\alwaysuseapproximateconstants}%
43 \NewDocumentEnvironment{usepreciseconstants}{}{\alwaysusepreciseconstants}%
```

mandi now has a key-value interface, implemented with `pgfkeys` and `pgfkeys`. There are two options: `units`<sup>P.8</sup>, with values `base`, `derived`, or `alternate` selects the default form of units `preciseconstants`<sup>P.8</sup>, with values `true` and `false`, selects precise numerical values for constants rather than approximate values.

First, define the keys. The key handlers require certain commands defined by the unit engine.

```
44 \newif\ifusingpreciseconstants
45 \pgfkeys{%
46   /mandi/options/.cd,
47   initial@setup/.style={%
48     /mandi/options/buffered@units/.initial=alternate,%}
49   },%
50   initial@setup,%  

51   preciseconstants/.is if=usingpreciseconstants,%  

52   units/.is choice,%  

53   units/.default=derived,%  

54   units/alternate/.style={/mandi/options/buffered@units=alternate},%  

55   units/base/.style={/mandi/options/buffered@units=base},%  

56   units/derived/.style={/mandi/options/buffered@units=derived},%  

57   .unknown/.code={%
58     \typeout{}%
59     \typeout{mandi: You used unknown option '\pgfkeyscurrentname'.}%
60   },%
61 }%
```

Process the options.

```
62 \ProcessPgfPackageOptions{/mandi/options}
```

Write a banner to the console showing the options in use.

```
63 \typeout{}%
64 \typeout{mandi: You are using mandi \mandiversion.}%
65 \typeout{mandi: This package requires LuaLaTeX.}%
66 \typeout{mandi: Loadtime options...}
```

Complete the banner by showing currently selected options. The value of the `units`<sup>P.8</sup> key is used in situ to set the default units.

```
67 \newcommand*\@mandi@do@setup}{%
68   \csname alwaysuse\pgfkeysvalueof{/mandi/options/buffered@units}units\endcsname%
69   \typeout{mandi: You will get \pgfkeysvalueof{/mandi/options/buffered@units}\space units.}%
70   \ifusingpreciseconstants
71     \alwaysusepreciseconstants
72     \typeout{mandi: You will get precise constants.}%
73   \else
74     \alwaysuseapproximateconstants
75     \typeout{mandi: You will get approximate constants.}%
76   \fi
77   \typeout{}%
78 }%
79 \mandi@do@setup
```

Define a setup command that overrides the loadtime options when called with new options. A new banner is written to the console.



```

136 \ExplSyntaxOn
137 \cs_new:Npn \__mandi_newscalarquantity:nnnn #1#2#3#4
138 {
139     \cs_new:cpx {#1} ##1 {\unit{##1}{\mandi@selectunits{#2}{#3}{#4}}}
140     \cs_new:cpx {#1value} ##1 {##1}
141     \cs_new:cpx {#1baseunits} ##1 {\unit{##1}{\mandi@selectbaseunits{#2}{#3}{#4}}}
142     \cs_new:cpx {#1derivedunits} ##1 {\unit{##1}{\mandi@selectderivedunits{#2}{#3}{#4}}}
143     \cs_new:cpx {#1alternateunits} ##1 {\unit{##1}{\mandi@selectalternateunits{#2}{#3}{#4}}}
144     \cs_new:cpx {#1onlybaseunits} {\mandi@selectbaseunits{#2}{#3}{#4}}
145     \cs_new:cpx {#1onlyderivedunits} {\mandi@selectderivedunits{#2}{#3}{#4}}
146     \cs_new:cpx {#1onlyalternateunits} {\mandi@selectalternateunits{#2}{#3}{#4}}
147 }
148 \NewDocumentCommand{\newscalarquantity}{ m m O{#2} O{#2} }
149 {
150     \__mandi_newscalarquantity:nnnn { #1 }{ #2 }{ #3 }{ #4 }
151 }
152 \ExplSyntaxOff

```

Redefining an existing scalar quantity.

```

153 \ExplSyntaxOn
154 \cs_new:Npn \__mandi_renewsscalarquantity:nnnn #1#2#3#4
155 {
156     \cs_set:cpx {#1} ##1 {\unit{##1}{\mandi@selectunits{#2}{#3}{#4}}}
157     \cs_set:cpx {#1value} ##1 {##1}
158     \cs_set:cpx {#1baseunits} ##1 {\unit{##1}{\mandi@selectbaseunits{#2}{#3}{#4}}}
159     \cs_set:cpx {#1derivedunits} ##1 {\unit{##1}{\mandi@selectderivedunits{#2}{#3}{#4}}}
160     \cs_set:cpx {#1alternateunits} ##1 {\unit{##1}{\mandi@selectalternateunits{#2}{#3}{#4}}}
161     \cs_set:cpx {#1onlybaseunits} {\mandi@selectbaseunits{#2}{#3}{#4}}
162     \cs_set:cpx {#1onlyderivedunits} {\mandi@selectderivedunits{#2}{#3}{#4}}
163     \cs_set:cpx {#1onlyalternateunits} {\mandi@selectalternateunits{#2}{#3}{#4}}
164 }
165 \NewDocumentCommand{\renewsscalarquantity}{ m m O{#2} O{#2} }
166 {
167     \__mandi_renewsscalarquantity:nnnn { #1 }{ #2 }{ #3 }{ #4 }
168 }
169 \ExplSyntaxOff

```

Defining a new vector quantity. Note that a corresponding scalar is also defined.

```

170 \ExplSyntaxOn
171 \cs_new:Npn \__mandi_newvectorquantity:nnnn #1#2#3#4
172 {
173     \__mandi_newscalarquantity:nnnn { #1 }{ #2 }{ #3 }{ #4 }
174     \cs_new:cpx {vector#1} ##1 {\unit{\mivector{##1}}{\mandi@selectunits{#2}{#3}{#4}}}
175     \cs_new:cpx {#1vector} ##1 {\unit{\mivector{##1}}{\mandi@selectunits{#2}{#3}{#4}}}
176     \cs_new:cpx {vector#1value} ##1 {\mivector{##1}}
177     \cs_new:cpx {#1vectorvalue} ##1 {\mivector{##1}}
178     \cs_new:cpx {vector#1baseunits} ##1 {\unit{\mivector{##1}}{\mandi@selectbaseunits{#2}{#3}{#4}}}
179     \cs_new:cpx {#1vectorbaseunits} ##1 {\unit{\mivector{##1}}{\mandi@selectbaseunits{#2}{#3}{#4}}}
180     \cs_new:cpx {vector#1derivedunits} ##1 {\unit{\mivector{##1}}{\mandi@selectderivedunits{#2}{#3}{#4}}}
181     \cs_new:cpx {#1vectorderivedunits} ##1 {\unit{\mivector{##1}}{\mandi@selectderivedunits{#2}{#3}{#4}}}
182     \cs_new:cpx {vector#1alternateunits} ##1 {\unit{\mivector{##1}}{\mandi@selectalternateunits{#2}{#3}{#4}}}
183     \cs_new:cpx {#1vectoralternateunits} ##1 {\unit{\mivector{##1}}{\mandi@selectalternateunits{#2}{#3}{#4}}}
184     \cs_new:cpx {vector#1onlybaseunits} {\mandi@selectbaseunits{#2}{#3}{#4}}
185     \cs_new:cpx {#1vectoronlybaseunits} {\mandi@selectbaseunits{#2}{#3}{#4}}
186     \cs_new:cpx {vector#1onlyderivedunits} {\mandi@selectderivedunits{#2}{#3}{#4}}
187     \cs_new:cpx {#1vectoronlyderivedunits} {\mandi@selectderivedunits{#2}{#3}{#4}}
188     \cs_new:cpx {vector#1onlyalternateunits} {\mandi@selectalternateunits{#2}{#3}{#4}}
189     \cs_new:cpx {#1vectoronlyalternateunits} {\mandi@selectalternateunits{#2}{#3}{#4}}
190 }

```

```

191 \NewDocumentCommand{\newvectorquantity}{ m m O{#2} O{#2} }
192   {
193     \__mandi_newvectorquantity:nnnn { #1 }{ #2 }{ #3 }{ #4 }
194   }
195 \ExplSyntaxOff

  Redefining an existing vector quantity. Note that a corresponding scalar is also redefined.

196 \ExplSyntaxOn
197 \cs_new:Npn \__mandi_renewvectorquantity:nnnn #1#2#3#4
198   {
199     \__mandi_renewscalarquantity:nnnn { #1 }{ #2 }{ #3 }{ #4 }
200     \cs_set:cpx {vector#1} ##1 {\unit{\mivector{##1}}{\mandi@selectunits{#2}{#3}{#4}}}
201     \cs_set:cpx {#1vector} ##1 {\unit{\mivector{##1}}{\mandi@selectunits{#2}{#3}{#4}}}
202     \cs_set:cpx {vector#1value} ##1 {\mivector{##1}}
203     \cs_set:cpx {#1vectorvalue} ##1 {\mivector{##1}}
204     \cs_set:cpx {vector#1baseunits} ##1 {\unit{\mivector{##1}}{\mandi@selectbaseunits{#2}{#3}{#4}}}
205     \cs_set:cpx {#1vectorbaseunits} ##1 {\unit{\mivector{##1}}{\mandi@selectbaseunits{#2}{#3}{#4}}}
206     \cs_set:cpx {vector#1derivedunits} ##1 {\unit{\mivector{##1}}{\mandi@selectderivedunits{#2}{#3}{#4}}}
207     \cs_set:cpx {#1vectorderivedunits} ##1 {\unit{\mivector{##1}}{\mandi@selectderivedunits{#2}{#3}{#4}}}
208     \cs_set:cpx {vector#1alternateunits} ##1 {\unit{\mivector{##1}}{\mandi@selectalternateunits{#2}{#3}{#4}}}
209     \cs_set:cpx {#1vectoralternateunits} ##1 {\unit{\mivector{##1}}{\mandi@selectalternateunits{#2}{#3}{#4}}}
210     \cs_set:cpx {vector#1onlybaseunits} {\mandi@selectbaseunits{#2}{#3}{#4}}
211     \cs_set:cpx {#1vectoronlybaseunits} {\mandi@selectbaseunits{#2}{#3}{#4}}
212     \cs_set:cpx {vector#1onlyderivedunits} {\mandi@selectderivedunits{#2}{#3}{#4}}
213     \cs_set:cpx {#1vectoronlyderivedunits} {\mandi@selectderivedunits{#2}{#3}{#4}}
214     \cs_set:cpx {vector#1onlyalternateunits} {\mandi@selectalternateunits{#2}{#3}{#4}}
215     \cs_set:cpx {#1vectoronlyalternateunits} {\mandi@selectalternateunits{#2}{#3}{#4}}
216   }
217 \NewDocumentCommand{\renewvectorquantity}{ m m O{#2} O{#2} }
218   {
219     \__mandi_renewvectorquantity:nnnn { #1 }{ #2 }{ #3 }{ #4 }
220   }
221 \ExplSyntaxOff

  Defining a new physical constant.

222 \ExplSyntaxOn
223 \cs_new:Npn \__mandi_newphysicalconstant:nnnnnnn #1#2#3#4#5#6#7
224   {
225     \cs_new:cpx {#1} {\unit{\mandi@selectprecision{#3}{#4}}{\mandi@selectunits{#5}{#6}{#7}}}
226     \cs_new:cpx {#1mathsymbol} {#2}
227     \cs_new:cpx {#1approximatevalue} {#3}
228     \cs_new:cpx {#1precisevalue} {#4}
229     \cs_new:cpx {#1baseunits}
230       {\unit{\mandi@selectprecision{#3}{#4}}{\mandi@selectbaseunits{#5}{#6}{#7}}}
231     \cs_new:cpx {#1derivedunits}
232       {\unit{\mandi@selectprecision{#3}{#4}}{\mandi@selectderivedunits{#5}{#6}{#7}}}
233     \cs_new:cpx {#1alternateunits}
234       {\unit{\mandi@selectprecision{#3}{#4}}{\mandi@selectalternateunits{#5}{#6}{#7}}}
235     \cs_new:cpx {#1onlybaseunits} {\mandi@selectbaseunits{#5}{#6}{#7}}
236     \cs_new:cpx {#1onlyderivedunits} {\mandi@selectderivedunits{#5}{#6}{#7}}
237     \cs_new:cpx {#1onlyalternateunits} {\mandi@selectalternateunits{#5}{#6}{#7}}
238   }
239 \NewDocumentCommand{\newphysicalconstant}{ m m m m m O{#5} O{#5} }
240   {
241     \__mandi_newphysicalconstant:nnnnnnn { #1 }{ #2 }{ #3 }{ #4 }{ #5 }{ #6 }{ #7 }
242   }
243 \ExplSyntaxOff

  Redefining an existing physical constant.

```

```

244 \ExplSyntaxOn
245 \cs_new:Npn \__mandi_renewphysicalconstant:nnnnnnn #1#2#3#4#5#6#7
246 {
247     \cs_set:cpn {#1} {\unit{\mandi@selectprecision{#3}{#4}}{\mandi@selectunits{#5}{#6}{#7}}}
248     \cs_set:cpn {#1mathsymbol} {#2}
249     \cs_set:cpn {#1approximatevalue} {#3}
250     \cs_set:cpn {#1precisevalue} {#4}
251     \cs_set:cpn {#1baseunits}
252         {\unit{\mandi@selectprecision{#3}{#4}}{\mandi@selectbaseunits{#5}{#6}{#7}}}
253     \cs_set:cpn {#1derivedunits}
254         {\unit{\mandi@selectprecision{#3}{#4}}{\mandi@selectderivedunits{#5}{#6}{#7}}}
255     \cs_set:cpn {#1alternateunits}
256         {\unit{\mandi@selectprecision{#3}{#4}}{\mandi@selectalternateunits{#5}{#6}{#7}}}
257     \cs_set:cpn {#1onlybaseunits} {\mandi@selectbaseunits{#5}{#6}{#7}}
258     \cs_set:cpn {#1onlyderivedunits} {\mandi@selectderivedunits{#5}{#6}{#7}}
259     \cs_set:cpn {#1onlyalternateunits} {\mandi@selectalternateunits{#5}{#6}{#7}}
260 }
261 \NewDocumentCommand{\renewphysicalconstant}{ m m m m m O{#5} O{#5} }
262 {
263     \__mandi_renewphysicalconstant:nnnnnnn { #1 }{ #2 }{ #3 }{ #4 }{ #5 }{ #6 }{ #7 }
264 }
265 \ExplSyntaxOff

```

Define every quantity we need in introductory physics, alphabetically for convenience. This is really the core feature of `mandi` that no other package offers. There are commands for quantities that have no dimensions or units, and these quantities are defined for semantic completeness.

```

266 \newvectorquantity{acceleration}%
267   {\meter\usk\second\totheinversetwo}%
268   [\newton\per\kilogram]%
269   [\meter\per\second\tothetwo]%
270 \newscalarquantity{amount}%
271   {\mole}%
272 \newvectorquantity{angularacceleration}%
273   {\radian\usk\second\totheinversetwo}%
274   [\radian\per\second\tothetwo]%
275   [\radian\per\second\tothetwo]%
276 \newscalarquantity{angularfrequency}%
277   {\radian\usk\second\inverse}%
278   [\radian\per\second]%
279   [\radian\per\second]%
280 %\ifmandi@rotadians
281 %  \newphysicalquantity{angularimpulse}%
282 %    {\meter\tothetwo\usk\kilogram\usk\second\inverse\usk\radian\inverse}%
283 %    [\joule\usk\second\per\radian]%
284 %    [\newton\usk\meter\usk\second\per\radian]%
285 %  \newphysicalquantity{angularmomentum}%
286 %    {\meter\tothetwo\usk\kilogram\usk\second\inverse\usk\radian\inverse}%
287 %    [\kilogram\usk\meter\tothetwo\per(\second\usk\radian)]%
288 %    [\newton\usk\meter\usk\second\per\radian]%
289 %\else
290 \newvectorquantity{angularimpulse}%
291   {\kilogram\usk\meter\tothetwo\usk\second\inverse}%
292   [\kilogram\usk\meter\tothetwo\per\second] % also \joule\usk\second
293   [\kilogram\usk\meter\tothetwo\per\second] % also \newton\usk\meter\usk\second
294 \newvectorquantity{angularmomentum}%
295   {\kilogram\usk\meter\tothetwo\usk\second\inverse}%
296   [\kilogram\usk\meter\tothetwo\per\second] % also \joule\usk\second
297   [\kilogram\usk\meter\tothetwo\per\second] % also \newton\usk\meter\usk\second
298 %\fi

```

```

299 \newvectorquantity{angularvelocity}%
300   {\radian\usk\second\inverse}%
301   [\radian\per\second]%
302   [\radian\per\second]%
303 \newsscalarquantity{area}%
304   {\meter\tothetwo}%
305 \newsscalarquantity{areachargedensity}%
306   {\ampere\usk\second\usk\meter\totheinversetwo}%
307   [\coulomb\per\meter\tothetwo]%
308   [\coulomb\per\meter\tothetwo]%
309 \newsscalarquantity{areamassdensity}%
310   {\kilogram\usk\meter\totheinversetwo}%
311   [\kilogram\per\meter\tothetwo]%
312   [\kilogram\per\meter\tothetwo]%
313 \newsscalarquantity{capacitance}%
314   {\ampere\tothetwo\usk\second\tothefour\usk\kilogram\inverse\usk\meter\totheinversetwo}%
315   [\farad]%
316   [\coulomb\per\volt] % also \coulomb\tothetwo\per\newton\usk\meter, \second\per\ohm
317 \newsscalarquantity{charge}%
318   {\ampere\usk\second}%
319   [\coulomb]%
320   [\coulomb] % also \farad\usk\volt
321 \newvectorquantity{cmagneticfield}%
322   {\kilogram\usk\meter\usk\ampere\inverse\usk\second\totheinversethree}%
323   [\newton\per\coulomb] % also \volt\per\meter
324   [\newton\per\coulomb]%
325 \newsscalarquantity{conductance}%
326   {\ampere\tothetwo\usk\second\tothethree\usk\kilogram\inverse\usk\meter\totheinversetwo}%
327   [\siemens]%
328   [\ampere\per\volt]%
329 \newsscalarquantity{conductivity}%
330   {\ampere\tothetwo\usk\second\tothethree\usk\kilogram\inverse\usk\meter\totheinversethree}%
331   [\siemens\per\meter]%
332   [\ampere\per\volt\usk\meter]%
333 \newsscalarquantity{conventionalcurrent}%
334   {\ampere}%
335   [\coulomb\per\second]%
336   [\ampere]%
337 \newsscalarquantity{current}%
338   {\ampere}%
339 \newsscalarquantity{currentdensity}%
340   {\ampere\usk\meter\totheinversetwo}%
341   [\coulomb\per\second\usk\meter\tothetwo]%
342   [\ampere\per\meter\tothetwo]%
343 \newsscalarquantity{dielectricconstant}%
344   {}%
345 \newvectorquantity{direction}%
346   {}%
347 \newvectorquantity{displacement}%
348   {\meter}%
349 \newsscalarquantity{duration}%
350   {\second}%
351 \newvectorquantity{electricdipolemoment}%
352   {\ampere\usk\second\usk\meter}%
353   [\coulomb\usk\meter]%
354   [\coulomb\usk\meter]%
355 \newvectorquantity{electricfield}%
356   {\kilogram\usk\meter\usk\ampere\inverse\usk\second\totheinversethree}%
357   [\volt\per\meter]%

```

```

358  [\newton\per\coulomb]%
359 \newscalarquantity{electricflux}%
360  {\kilogram\usk\meter\tothethree\usk\ampere\inverse\usk\second\totheinversethtree}%
361  [\volt\usk\meter]%
362  [\newton\usk\meter\tothetwo\per\coulomb]%
363 \newscalarquantity{electricpotential}%
364  {\kilogram\usk\meter\tothetwo\usk\ampere\inverse\usk\second\totheinversethtree}%
365  [\volt]% % also \joule\per\coulomb
366  [\volt]%
367 \newscalarquantity{electricpotentiaifference}%
368  {\kilogram\usk\meter\tothetwo\usk\ampere\inverse\usk\second\totheinversethtree}%
369  [\volt]% % also \joule\per\coulomb
370  [\volt]%
371 \newscalarquantity{electroncurrent}%
372  {\second\inverse}%
373  [\ensuremath{\text{\textnormal{\textsf{symup}}}\{e\}}\per\second]%
374  [\ensuremath{\text{\textnormal{\textsf{symup}}}\{e\}}\per\second]%
375 \newscalarquantity{emf}%
376  {\kilogram\usk\meter\tothetwo\usk\ampere\inverse\usk\second\totheinversethtree}%
377  [\volt]% % also \joule\per\coulomb
378  [\volt]%
379 \newscalarquantity{energy}%
380  {\kilogram\usk\meter\tothetwo\usk\second\totheinversetwo}%
381  [\joule]% % also \newton\usk\meter
382  [\joule]%
383 \newscalarquantity{energyinev}%
384  {\electronvolt}%
385 \newscalarquantity{energyinkev}%
386  {\kilolectronvolt}%
387 \newscalarquantity{energyinmev}%
388  {\megaelectronvolt}%
389 \newscalarquantity{energydensity}%
390  {\kilogram\usk\meter\inverse\usk\second\totheinversetwo}%
391  [\joule\per\meter\tothethree]%
392  [\joule\per\meter\tothethree]%
393 \newscalarquantity{energyflux}%
394  {\kilogram\usk\second\totheinversethtree}%
395  [\watt\per\meter\tothetwo]%
396  [\watt\per\meter\tothetwo]%
397 \newscalarquantity{entropy}%
398  {\kilogram\usk\meter\tothetwo\usk\second\totheinversetwo\usk\kelvin\inverse}%
399  [\joule\per\kelvin]%
400  [\joule\per\kelvin]%
401 \newvectorquantity{force}%
402  {\kilogram\usk\meter\usk\second\totheinversetwo}%
403  [\newton]%
404  [\newton]% % also \kilogram\usk\meter\per\second\tothetwo
405 \newscalarquantity{frequency}%
406  {\second\inverse}%
407  [\hertz]%
408  [\hertz]%
409 \newvectorquantity{gravitationalfield}%
410  {\meter\usk\second\totheinversetwo}%
411  [\newton\per\kilogram]%
412  [\newton\per\kilogram]%
413 \newscalarquantity{gravitationalpotential}%
414  {\meter\tothetwo\usk\second\totheinversetwo}%
415  [\joule\per\kilogram]%
416  [\joule\per\kilogram]%

```

```

417 \newscalarquantity{gravitationalpotentialdifference}%
418   {\meter\tothetwo\usk\second\totheinversetwo}%
419   [\joule\per\kilogram]%
420   [\joule\per\kilogram]%
421 \newvectorquantity{impulse}%
422   {\kilogram\usk\meter\usk\second\inverse}%
423   [\newton\usk\second]%
424   [\newton\usk\second]%
425 \newscalarquantity{indexofrefraction}%
426   {}%
427 \newscalarquantity{inductance}%
428   {\kilogram\usk\meter\tothetwo\usk\ampere\totheinversetwo\usk\second\totheinversetwo}%
429   [\henry]%
430   [\volt\usk\second\per\ampere] % also \square\meter\usk\kilogram\per\coulomb\tothetwo, \Wb\per\ampere
431 \newscalarquantity{linearchargedensity}%
432   {\ampere\usk\second\usk\meter\inverse}%
433   [\coulomb\per\meter]%
434   [\coulomb\per\meter]%
435 \newscalarquantity{linearmassdensity}%
436   {\kilogram\usk\meter\inverse}%
437   [\kilogram\per\meter]%
438   [\kilogram\per\meter]%

439 \newscalarquantity{lorentzfactor}%
440   {}%
441 \newscalarquantity{luminousintensity}%
442   {\candela}%
443 \newscalarquantity{magneticcharge}%
444   {\ampere\usk\meter} % There is another convention. Be careful!
445 \newvectorquantity{magneticdipolemoment}%
446   {\ampere\usk\meter\tothetwo}%
447   [\ampere\usk\meter\tothetwo]%
448   [\joule\per\tesla]%
449 \newvectorquantity{magneticfield}%
450   {\kilogram\usk\ampere\inverse\usk\second\totheinversetwo}%
451   [\newton\per\ampere\usk\meter] % also \Wb\per\meter\tothetwo
452   [\tesla]%
453 \newscalarquantity{magneticflux}%
454   {\kilogram\usk\meter\tothetwo\usk\ampere\inverse\usk\second\totheinversetwo}%
455   [\tesla\usk\meter\tothetwo]%
456   [\volt\usk\second] % also \Wb and \joule\per\ampere
457 \newscalarquantity{mass}%
458   {\kilogram}%
459 \newscalarquantity{mobility}%
460   {\kilogram\usk\meter\tothetwo\usk\ampere\inverse\usk\second\totheinversefour}%
461   [\meter\tothetwo\per\volt\usk\second]%
462   [\coulomb\usk\meter\per\newton\usk\second]%
463 \newscalarquantity{momentofinertia}%
464   {\kilogram\usk\meter\tothetwo}%
465   [\joule\usk\second\tothetwo]%
466   [\kilogram\usk\meter\tothetwo]%
467 \newvectorquantity{momentum}%
468   {\kilogram\usk\meter\usk\second\inverse}%
469   [\kilogram\usk\meter\per\second]%
470   [\kilogram\usk\meter\per\second]%
471 \newvectorquantity{momentumflux}%
472   {\kilogram\usk\meter\inverse\usk\second\totheinversetwo}%
473   [\newton\per\meter\tothetwo]%
474   [\newton\per\meter\tothetwo]%

```

```

475 \newscalarquantity{numberdensity}%
476   {\meter\tothe^{-3}}%
477   [\per\meter\tothe{3}]%
478   [\per\meter\tothe{3}]%
479 \newscalarquantity{permeability}%
480   {\kilogram\usk\meter\usk\ampere\tothe^{-2}\usk\second\tothe^{-2}}%
481   [\henry\per\meter]%
482   [\tesla\usk\meter\per\ampere]%
483 \newscalarquantity{permittivity}%
484   {\ampere\tothe{2}\usk\second\tothe{4}\usk\kilogram\inverse\usk\meter\tothe{-3}}%
485   [\farad\per\meter]%
486   [\coulomb\tothe{2}\per\newton\usk\meter\tothe{2}]%
487 \newscalarquantity{planeangle}%
488   {\meter\usk\meter\inverse}%
489   [\radian]%
490   [\radian]%
491 \newscalarquantity{polarizability}%
492   {\ampere\tothe{2}\usk\second\tothe{4}\usk\kilogram\inverse}%
493   [\coulomb\usk\meter\tothe{2}\per\volt]%
494   [\coulomb\tothe{2}\usk\meter\per\newton]%
495 \newscalarquantity{power}%
496   {\kilogram\usk\meter\tothe{2}\usk\second\tothe{-3}}%
497   [\watt]%
498   [\joule\per\second]%
499 \newvectorquantity{poynting}%
500   {\kilogram\usk\second\tothe{-3}}%
501   [\watt\per\meter\tothe{2}]%
502   [\watt\per\meter\tothe{2}]%
503 \newscalarquantity{pressure}%
504   {\kilogram\usk\meter\inverse\usk\second\tothe{-2}}%
505   [\pascal]%
506   [\newton\per\meter\tothe{2}]%
507 \newscalarquantity{relativepermeability}%
508   {}%
509 \newscalarquantity{relativepermittivity}%
510   {}%
511 \newscalarquantity{resistance}%
512   {\kilogram\usk\meter\tothe{2}\usk\ampere\tothe{-2}\usk\second\tothe{-3}}%
513   [\ohm] % also \volt\per\ampere
514   [\ohm]%
515 \newscalarquantity{resistivity}%
516   {\kilogram\usk\meter\tothe{3}\usk\ampere\tothe{-2}\usk\second\tothe{-3}}%
517   [\ohm\usk\meter]%
518   [\volt\usk\meter\per\ampere]%
519 \newscalarquantity{solidangle}%
520   {\meter\tothe{2}\usk\meter\tothe{-2}}%
521   [\steradian]%
522   [\steradian]%
523 \newscalarquantity{specificheatcapacity}%
524   {\meter\tothe{2}\usk\second\tothe{-2}\usk\kelvin\inverse}%
525   [\joule\per\kelvin\usk\kilogram]%
526   [\joule\per\kelvin\usk\kilogram]%
527 \newscalarquantity{springstiffness}%
528   {\kilogram\usk\second\tothe{-2}}%
529   [\newton\per\meter]%
530   [\newton\per\meter]%
531 \newscalarquantity{springstretch} % This is really just a displacement.
532   {\meter}%
533 \newscalarquantity{stress}%

```

```

534  {\kilogram\usk\meter\inverse\usk\second\totheinversetwo}%
535  [\pascal]%
536  [\newton\per\meter\tothetwo]%
537 \newscalarquantity{strain}%
538 {}%
539 \newscalarquantity{temperature}%
540  {\kelvin}%
541 %\ifmandi@rotadians
542 %  \newphysicalquantity{torque}%
543 %  {\kilogram\usk\meter\tothetwo\usk\second\totheinversetwo\usk\radian\inverse}%
544 %  [\newton\usk\meter\per\radian]%
545 %  [\newton\usk\meter\per\radian]%
546 %\else
547 \newvectorquantity{torque}%
548  {\kilogram\usk\meter\tothetwo\usk\second\totheinversetwo}%
549  [\newton\usk\meter]%
550  [\newton\usk\meter]%
551 %\fi
552 \newvectorquantity{velocity}%
553  {\meter\usk\second\inverse}%
554  [\meter\per\second]%
555  [\meter\per\second]%
556 \newvectorquantity{velocityc}%
557  {\lightspeed}%
558  [\lightspeed]%
559  [\lightspeed]%
560 \newscalarquantity{volume}%
561  {\meter\tothethree}%
562 \newscalarquantity{volumechargedensity}%
563  {\ampere\usk\second\per\meter\totheinversethree}%
564  [\coulomb\per\meter\tothethree]%
565  [\coulomb\per\meter\tothethree]%
566 \newscalarquantity{volumemassdensity}%
567  {\kilogram\usk\meter\totheinversethree}%
568  [\kilogram\per\meter\tothethree]%
569  [\kilogram\per\meter\tothethree]%
570 \newscalarquantity{wavelength} % This is really just a displacement.
571  {\meter}%
572 \newvectorquantity{wavenumber}%
573  {\meter\inverse}%
574  [\per\meter]%
575  [\per\meter]%
576 \newscalarquantity{work}%
577  {\kilogram\usk\meter\tothetwo\usk\second\totheinversetwo}%
578  [\joule] % also \newton\usk\meter but discouraged
579  [\joule]%
580 \newscalarquantity{youngsmodulus} % This is really just a stress.
581  {\kilogram\usk\meter\inverse\usk\second\totheinversetwo}%
582  [\pascal]%
583  [\newton\per\meter\tothetwo]%

```

We need a better glyph for Planck's constant over  $2\pi$ .

```

584 \AtBeginDocument{%
585   \DeclareRobustCommand{\hbar}{\mathpalette\hbar@\relax\symup{h}}}%
586 }%
587 \newcommand*{\hbar@}[2]{%
588   \makebox[0pt][l]{\raisebox{-0.07\height}{\(\m@th#1\mkern-2mu\mathchar"AF\)}}}%
589   % optional line to make the bar thicker; must use -0.11
590   \makebox[0pt][l]{\raisebox{-0.11\height}{\(\m@th#1\mkern-2mu\mathchar"AF\)}}}%

```

591 }%

Define physical constants for introductory physics, again alphabetically for convenience.

```
592 \newphysicalconstant{avogadro}%
593   {\symup{N_A}}%
594   {6\times10^{-23}}{6.02214076\times10^{-23}}% % exact 2019 value
595   {\mole\inverse}%
596   [\per\mole]%
597   [\per\mole]%
598 \newphysicalconstant{biotsavartconstant} % alias for \mzofp
599   {\symup{\frac{\mu_0}{4\pi}}}}%
600   {10^{-7}}{10^{-7}}%
601   {\kilogram\usk\meter\usk\ampere\totheinversetwo\usk\second\totheinversetwo}%
602   [\henry\per\meter]%
603   [\tesla\usk\meter\per\ampere]%
604 \newphysicalconstant{bohradius}%
605   {\symup{a_o}}%
606   {5.3\times10^{-11}}{5.29177210903\times10^{-11}}%
607   {\meter}%
608 \newphysicalconstant{boltzmann}%
609   {\symup{k_B}}%
610   {1.4\times10^{-23}}{1.380649\times10^{-23}}% % exact 2019 value
611   {\kilogram\usk\meter\tothetwo\usk\second\totheinversetwo\usk\kelvin\inverse}%
612   [\joule\per\kelvin]%
613   [\joule\per\kelvin]%
614 \newphysicalconstant{coulombconstant} % alias for \oofpez
615   {\symup{\frac{1}{4\pi\epsilon_0}}}}%
616   {9\times10^{-9}}{8.9875517923\times10^{-9}}%
617   {\kilogram\usk\meter\tothethree\usk\ampere\totheinversetwo\usk\second\totheinversefour}%
618   [\meter\per\farad]%
619   [\newton\usk\meter\tothetwo\per\coulomb\tothetwo]%
620 \newphysicalconstant{earthmass}%
621   {\symup{M_Earth}}%
622   {6.0\times10^{24}}{5.9722\times10^{24}}%
623   {\kilogram}%
624 \newphysicalconstant{earthmoondistance}%
625   {\symup{d_EM}}%
626   {3.8\times10^8}{3.81550\times10^8}%
627   {\meter}%
628 \newphysicalconstant{earthradius}%
629   {\symup{R_Earth}}%
630   {6.4\times10^6}{6.3781\times10^6}%
631   {\meter}%
632 \newphysicalconstant{earthsundistance}%
633   {\symup{d_ES}}%
634   {1.5\times10^{11}}{1.496\times10^{11}}%
635   {\meter}%
636 \newphysicalconstant{electroncharge}%
637   {\symup{q_e}}%
638   {-\elementarychargeapproximatevalue}{-\elementarychargeprecisevalue}%
639   {\ampere\usk\second}%
640   [\coulomb]%
641   [\coulomb]%
642 \newphysicalconstant{electronCharge}%
643   {\symup{Q_e}}%
644   {-\elementarychargeapproximatevalue}{-\elementarychargeprecisevalue}%
645   {\ampere\usk\second}%
646   [\coulomb]%
647   [\coulomb]
```

```

648 \newphysicalconstant{electronmass}%
649   {\symup{m_e}}%
650   {9.1\times10^{-31}}{9.1093837015\times10^{-31}}%
651   {\kilogram}%
652 \newphysicalconstant{elementarycharge}%
653   {\symup{e}}%
654   {1.6\times10^{-19}}{1.602176634\times10^{-19}}% exact 2019 value
655   {\ampere\usk\second}%
656   [\coulomb]%
657   [\coulomb]%
658 \newphysicalconstant{finestructure}%
659   {\symup{\alpha}}%
660   {\frac{1}{137}}{7.2973525693\times10^{-3}}%
661   {}%
662 \newphysicalconstant{hydrogenmass}%
663   {\symup{m_H}}%
664   {1.7\times10^{-27}}{1.6737236\times10^{-27}}%
665   {\kilogram}%
666 \newphysicalconstant{moonearthdistance}%
667   {\symup{d_ME}}%
668   {3.8\times10^8}{3.81550\times10^8}%
669   {\meter}%
670 \newphysicalconstant{moonmass}%
671   {\symup{M_Moon}}%
672   {7.3\times10^{22}}{7.342\times10^{22}}%
673   {\kilogram}%
674 \newphysicalconstant{moonradius}%
675   {\symup{R_Moon}}%
676   {1.7\times10^6}{1.7371\times10^6}%
677   {\meter}%
678 \newphysicalconstant{mzofp}%
679   {\symup{\frac{\mu_0}{4\pi}}}%
680   {10^{-7}}{10^{-7}}%
681   {\kilogram\usk\meter\usk\ampere\totheinversetwo\usk\second\totheinversetwo}%
682   [\henry\per\meter]%
683   [\tesla\usk\meter\per\ampere]%
684 \newphysicalconstant{neutronmass}%
685   {\symup{m_n}}%
686   {1.7\times10^{-27}}{1.67492749804\times10^{-27}}%
687   {\kilogram}%
688 \newphysicalconstant{oofpez}%
689   {\symup{\frac{1}{4\pi\epsilon_0}}}%
690   {9\times10^{-9}}{8.9875517923\times10^{-9}}%
691   {\kilogram\usk\meter\tothethree\usk\ampere\totheinversetwo\usk\second\totheinversefour}%
692   [\meter\per\farad]%
693   [\newton\usk\meter\tothetwo\per\coulomb\tothetwo]%
694 \newphysicalconstant{oofpezcs}%
695   {\symup{\frac{1}{4\pi\epsilon_0 c^2}}}%
696   {10^{-7}}{10^{-7}}%
697   {\kilogram\usk\meter\usk\ampere\totheinversetwo\usk\second\totheinversetwo}%
698   [\tesla\usk\meter\tothetwo]%
699   [\newton\usk\second\tothetwo\per\coulomb\tothetwo]%
700 \newphysicalconstant{planck}%
701   {\symup{h}}%
702   {6.6\times10^{-34}}{6.62607015\times10^{-34}}% exact 2019 value
703   {\kilogram\usk\meter\tothetwo\usk\second\inverse}%
704   [\joule\usk\second]%
705   [\joule\usk\second]%

```

See <https://tex.stackexchange.com/a/448565/218142>.

```
706 \newphysicalconstant{planckbar}%
707   {\hbar}%
708   {1.1\times10^{-34}}{1.054571817\times10^{-34}}%
709   {\kilogram\usk\meter\tothetwo\usk\second\inverse}%
710   [\joule\usk\second]%
711   [\joule\usk\second]
712 \newphysicalconstant{planckc}%
713   {\symup{hc}}%
714   {2.0\times10^{-25}}{1.98644586\times10^{-25}}%
715   {\kilogram\usk\meter\tothethree\usk\second\totheinversetwo}%
716   [\joule\usk\meter]%
717   [\joule\usk\meter]%
718 \newphysicalconstant{protoncharge}%
719   {\symup{q_p}}%
720   {+\elementarychargeapproximatevalue}{+\elementarychargeprecisevalue}%
721   {\ampere\usk\second}%
722   [\coulomb]%
723   [\coulomb]%
724 \newphysicalconstant{protonCharge}%
725   {\symup{Q_p}}%
726   {+\elementarychargeapproximatevalue}{+\elementarychargeprecisevalue}%
727   {\ampere\usk\second}%
728   [\coulomb]%
729   [\coulomb]%
730 \newphysicalconstant{protonmass}%
731   {\symup{m_p}}%
732   {1.7\times10^{-27}}{1.672621898\times10^{-27}}%
733   {\kilogram}%
734 \newphysicalconstant{rydberg}%
735   {\symup{R_{\infty}}}%
736   {1.1\times10^{7}}{1.0973731568160\times10^{7}}%
737   {\meter\inverse}%
738 \newphysicalconstant{speedoflight}%
739   {\symup{c}}%
740   {3\times10^{8}}{2.99792458\times10^{8}}% exact value
741   {\meter\usk\second\inverse}%
742   [\meter\per\second]%
743   [\meter\per\second]
744 \newphysicalconstant{stefanboltzmann}%
745   {\symup{\sigma}}%
746   {5.7\times10^{-8}}{5.670374\times10^{-8}}%
747   {\kilogram\usk\second\totheinversethree\usk\kelvin\totheinversefour}%
748   [\watt\per\meter\tothetwo\usk\kelvin\tothefour]%
749   [\watt\per\meter\tothetwo\usk\kelvin\tothefour]
750 \newphysicalconstant{sunearthdistance}%
751   {\symup{d_SE}}%
752   {1.5\times10^{11}}{1.496\times10^{11}}%
753   {\meter}%
754 \newphysicalconstant{sunmass}%
755   {\symup{M_Sun}}%
756   {2.0\times10^{30}}{1.98855\times10^{30}}%
757   {\kilogram}%
758 \newphysicalconstant{sunradius}%
759   {\symup{R_Sun}}%
760   {7.0\times10^{8}}{6.957\times10^{8}}%
761   {\meter}%
762 \newphysicalconstant{surfacegravfield}%
```

```

763 {\symup{g}}%
764 {9.8}{9.807}%
765 {\meter\usk\second\totheinversetwo}%
766 [\newton\per\kilogram]%
767 [\newton\per\kilogram]%
768 \newphysicalconstant{universalgrav}%
769 {\symup{G}}%
770 {6.7\times10^{-11}}{6.67430\times10^{-11}}%
771 {\meter\tothethree\usk\kilogram\inverse\usk\second\totheinversetwo}%
772 [\newton\usk\meter\tothetwo\per\kilogram\tothetwo] % also \joule\usk\meter\per\kilogram\tothetwo
773 [\newton\usk\meter\tothetwo\per\kilogram\tothetwo]%
774 \newphysicalconstant{vacuumpermability}%
775 {\symup{\mu_o}}%
776 {4\pi\times10^{-7}}{4\pi\times10^{-7}}% as of 2018 no longer 4\pi\times10^{-7}
777 {\kilogram\usk\meter\usk\ampere\totheinversetwo\usk\second\totheinversetwo}%
778 [\henry\per\meter]%
779 [\tesla\usk\meter\per\ampere]%
780 \newphysicalconstant{vacuumpermittivity}%
781 {\symup{\epsilon_0}}%
782 {9\times10^{-12}}{8.854187817\times10^{-12}}%
783 {\ampere\tothetwo\usk\second\tothefour\usk\kilogram\inverse\usk\meter\totheinversethree}%
784 [\farad\per\meter]%
785 [\coulomb\tothetwo\per\newton\usk\meter\tothetwo]%

```

Diagnostic commands to provide sanity checks on commands that represent physical quantities and constants.

```

786 \ExplSyntaxOn
787 \NewDocumentCommand{\@aux}{ m }
788 {
789     \use:c { #1 }
790 }
791 \NewDocumentCommand{\@auy}{ m }
792 {
793     \normalfont\ttfamily\token_to_str:c { #1 }
794 }
795 \ExplSyntaxOff
796 \newcolumntype{M}{>{\()p{0.25\linewidth}<{\()}}
797 \NewDocumentCommand{\checkquantity}{ m }
798 {%
799     \begin{center}
800         \begin{tabular}{MMM}
801             \textbf{command} & \multicolumn{2}{l}{\@auy{#1}} & \tabularnewline
802             \text{\textbf{base}} & \text{\textbf{derived}} & \text{\textbf{alternate}} & \tabularnewline
803             \@aux{#1onlybaseunits} & \@aux{#1onlyderivedunits} & \@aux{#1onlyalternateunits} & \tabularnewline
804         \end{tabular}
805     \end{center}
806 }%
807 \NewDocumentCommand{\checkconstant}{ m }
808 {%
809     \begin{center}
810         \begin{tabular}{MMM}
811             \textbf{command} & \multicolumn{2}{l}{\@auy{#1}} & \tabularnewline
812             \text{\textbf{symbol}} & \text{\textbf{approximate}} & \text{\textbf{precise}} & \tabularnewline
813             \@aux{#1mathsymbol} & \@aux{#1approximatevalue} & \@aux{#1precisevalue} & \tabularnewline
814             \text{\textbf{base}} & \text{\textbf{derived}} & \text{\textbf{alternate}} & \tabularnewline
815             \@aux{#1onlybaseunits} & \@aux{#1onlyderivedunits} & \@aux{#1onlyalternateunits} & \tabularnewline
816         \end{tabular}
817     \end{center}
818 }%

```

`\mivector`<sup>P.34</sup> is a workhorse command.

See <https://tex.stackexchange.com/a/39054/218142>.

```
819 \ExplSyntaxOn
820 \NewDocumentCommand{\mivector}{ O{,} m o }
821   {
822     \__mandi_vector:nn { #1 } { #2 }
823     \IfValueT{#3}{\, #3}
824   }
825 \seq_new:N \l__mandi_list_seq
826 \cs_new_protected:Npn \__mandi_vector:nn #1#2
827   {
828     \seq_set_split:Nnn \l__mandi_list_seq { , } { #2 }
829     \int_compare:nT { \seq_count:N \l__mandi_list_seq = 1 }
830     {
831       \msg_new:nnnn { mandi } { onecomponent }
832       {
833         More~than~one~component~expected.           \iow_newline:
834         You~provided~one~component~to~a~command \iow_newline:
835         that~expects~a~vector.~Either~you~don't \iow_newline:
836         need~a~vector~here~or~you~didn't~supply \iow_newline:
837         all~the~components.
838       }
839       {
840         Decide~whether~or~not~you~really~need~a~vector~command~here. \iow_newline:
841         \msg_see_documentation_text:n { mandi }
842       }
843       \msg_fatal:nn { mandi } { onecomponent }
844     }
845   }
846   \left\langle
847     \seq_use:Nnnn \l__mandi_list_seq { #1 } { #1 } { #1 }
848   \right\rangle
849 }
850 \ExplSyntaxOff
```

## 5 The `mandistudent` Package

`mandi` comes with an accessory package `mandistudent`, which provides a collection of commands physics students can use for writing problem solutions. This package focuses on the most frequently needed tools. These commands should always be used in math mode. Note that `mandistudent` requires, and loads, `mandi` but `mandi` doesn't require, and doesn't load, `mandistudent`.

Load `mandistudent` as you would any package in your preamble. There are no package options.

```
\usepackage{mandistudent}
```

`\mandistudentversion`

Typesets the current version and build date.

```
The version is \mandistudentversion\ and is a stable build.
```

```
The version is v3.1.0 dated 2022-01-27 and is a stable build.
```

### 5.1 Traditional Vector Notation

U 2021-09-18  
U 2021-09-18

`\vec{\symbol}{[\label]}`  
`\vec*{\symbol}{[\label]}`

(use this variant for boldface notation)  
(use this variant for arrow notation)

Powerful and intelligent command for symbolic vector notation. The mandatory argument is the symbol for the vector quantity. The optional label(s) consists of superscripts and/or subscripts and can be mathematical or textual in nature. If textual, be sure to wrap them in `\symup{...}` for proper typesetting. The starred variant gives arrow notation whereas without the star you get boldface notation. Subscript and superscript labels can be arbitrarily mixed, and order doesn't matter. This command redefines the default L<sup>A</sup>T<sub>E</sub>X 2 <sub>$\varepsilon$</sub>  `\vec` command.

<code>\(\vec{p}\)</code>	$p$
<code>\(\vec{p}_2\)</code>	$p_2$
<code>\(\vec{p}^{\text{ball}}\)</code>	$p^{\text{ball}}$
<code>\(\vec{p}_{\text{final}}\)</code>	$p_{\text{final}}$
<code>\(\vec{p}^{\text{final}}\)</code>	$p^{\text{final}}$
<code>\(\vec{p}^{\text{ball}}_{\text{final}}\)</code>	$p^{\text{ball}}_{\text{final}}$
<code>\(\vec{p}^{\text{final}}_{\text{ball}}\)</code>	$p^{\text{final}}_{\text{ball}}$
<code>\(\vec{p}\_2\)</code>	$\overrightarrow{p}$
<code>\(\vec{p}^{\text{ball}}\)</code>	$\overrightarrow{p}^{\text{ball}}$
<code>\(\vec{p}_{\text{final}}\)</code>	$\overrightarrow{p}_{\text{final}}$
<code>\(\vec{p}^{\text{final}}\)</code>	$\overrightarrow{p}^{\text{final}}$
<code>\(\vec{p}^{\text{final}}_{\text{ball}}\)</code>	$\overrightarrow{p}^{\text{final}}_{\text{ball}}$
<code>\(\vec{p}^{\text{ball}}_{\text{final}}\)</code>	$\overrightarrow{p}_{\text{final}}^{\text{ball}}$

U 2021-09-18  
U 2021-09-18

`\dirvec{<symbol>}[<labels>]`  
`\dirvec*{<symbol>}[<labels>]`

(use this variant for boldface notation)  
(use this variant for arrow notation)

Powerful and intelligent command for typesetting the direction of a vector. The options are the same as those for `\vec`.

<code>\(\dirvec{p}\)</code>	$\hat{\mathbf{p}}$
<code>\(\dirvec{p}_2\)</code>	$\hat{\mathbf{p}}_2$
<code>\(\dirvec{p}\`{\text{y}}{\text{mup}{ball}}\)</code>	$\hat{\mathbf{p}}_{\text{ball}}$
<code>\(\dirvec{p}\`{\text{y}}{\text{mup}{final}}\)</code>	$\hat{\mathbf{p}}_{\text{final}}$
<code>\(\dirvec{p}\`{\text{y}}{\text{mup}{ball}}\`{\text{y}}{\text{mup}{final}}\)</code>	$\hat{\mathbf{p}}_{\text{final}}^{\text{final}}$
<code>\(\dirvec{p}\`{\text{y}}{\text{mup}{final}}\`{\text{y}}{\text{mup}{ball}}\)</code>	$\hat{\mathbf{p}}_{\text{ball}}^{\text{final}}$
<code>\(\dirvec*{p}\)</code>	$\hat{\mathbf{p}}$
<code>\(\dirvec*{p}_2\)</code>	$\hat{\mathbf{p}}_2$
<code>\(\dirvec*{p}\`{\text{y}}{\text{mup}{ball}}\)</code>	$\hat{\mathbf{p}}_{\text{ball}}$
<code>\(\dirvec*{p}\`{\text{y}}{\text{mup}{final}}\)</code>	$\hat{\mathbf{p}}_{\text{final}}$
<code>\(\dirvec*{p}\`{\text{y}}{\text{mup}{ball}}\`{\text{y}}{\text{mup}{final}}\)</code>	$\hat{\mathbf{p}}_{\text{final}}^{\text{ball}}$
<code>\(\dirvec*{p}\`{\text{y}}{\text{mup}{final}}\`{\text{y}}{\text{mup}{ball}}\)</code>	$\hat{\mathbf{p}}_{\text{ball}}^{\text{final}}$

`\zerovec`  
`\zerovec*`

(use this variant for boldface notation)  
(use this variant for arrow notation)

Command for typesetting the zero vector. The starred variant gives arrow notation. Without the star you get boldface notation.

<code>\(\zerovec\)</code> \\	$\mathbf{0}$
<code>\(\zerovec*\)</code>	$\overrightarrow{0}$

`\changein`

Semantic alias for `\Delta`.

<code>\(\changein{t}\)</code> \\	$\Delta t$
<code>\(\changein{p}\)</code>	$\Delta p$

<code>\doublebars[&lt;size&gt;]{&lt;quantity&gt;}</code>	(double bars)
<code>\doublebars*[&lt;size&gt;]{&lt;quantity&gt;}</code>	(double bars for fractions)
<code>\singlebars[&lt;size&gt;]{&lt;quantity&gt;}</code>	(single bars)
<code>\singlebars*[&lt;size&gt;]{&lt;quantity&gt;}</code>	(single bars for fractions)
<code>\anglebrackets[&lt;size&gt;]{&lt;quantity&gt;}</code>	(angle brackets)
<code>\anglebrackets*[&lt;size&gt;]{&lt;quantity&gt;}</code>	(angle brackets for fractions)
<code>\parentheses[&lt;size&gt;]{&lt;quantity&gt;}</code>	(parentheses)
<code>\parentheses*[&lt;size&gt;]{&lt;quantity&gt;}</code>	(parentheses for fractions)

N 2021-02-21  
N 2021-02-21

N 2021-02-21  
 N 2021-02-21  
 N 2021-02-21  
 N 2021-02-21

<code>\squarebrackets[<i>size</i>]{<i>quantity</i>}</code>	(square brackets)
<code>\squarebrackets*[<i>size</i>]{<i>quantity</i>}</code>	(square brackets for fractions)
<code>\curlybraces[<i>size</i>]{<i>quantity</i>}</code>	(curly braces)
<code>\curlybraces*[<i>size</i>]{<i>quantity</i>}</code>	(curly braces for fractions)

If no argument is given, a placeholder is provided. Sizers like `\big`, `\Big`, `\bigg`, and `\Bigg` can be optionally specified. Beginners are encouraged not to use them. See the [mathtools](#) package documentation for details.

```
\[ \doublebars{} \]
\[ \doublebars{\vec{a}} \]
\[ \doublebars*{\frac{\vec{a}}{3}} \]
\[ \doublebars[\Bigg]{\frac{\vec{a}}{3}} \]
```

$$\| \cdot \|$$

$$\| a \|$$

$$\left\| \frac{a}{3} \right\|$$

$$\left\| \frac{a}{3} \right\|$$

```
\[ \singlebars{} \]
\[ \singlebars{x} \]
\[ \singlebars*{\frac{x}{3}} \]
\[ \singlebars[\Bigg]{\frac{x}{3}} \]
```

$$| \cdot |$$

$$| x |$$

$$\left| \frac{x}{3} \right|$$

$$\left| \frac{x}{3} \right|$$

```
\[ \anglebrackets{} \]
\[ \anglebrackets{\vec{a}} \]
\[ \anglebrackets*{\frac{\vec{a}}{3}} \]
\[ \anglebrackets[\Bigg]{\frac{\vec{a}}{3}} \]
```

$$\langle \cdot \rangle$$

$$\langle a \rangle$$

$$\left\langle \frac{a}{3} \right\rangle$$

$$\left\langle \frac{a}{3} \right\rangle$$

```
\[ \parentheses{} \]
\[ \parentheses{x} \]
\[ \parentheses*{\frac{x}{3}} \]
\[ \parentheses[\Bigg]{\frac{x}{3}} \]
```

$$\begin{aligned}(\cdot) \\ (x) \\ \left(\frac{x}{3}\right) \\ \left(\frac{x}{3}\right)\end{aligned}$$

```
\[ \squarebrackets{} \]
\[ \squarebrackets{x} \]
\[ \squarebrackets*{\frac{x}{3}} \]
\[ \squarebrackets[\Bigg]{\frac{x}{3}} \]
```

$$\begin{aligned}[\cdot] \\ [x] \\ \left[\frac{x}{3}\right] \\ \left[\frac{x}{3}\right]\end{aligned}$$

```
\[ \curlybraces{} \]
\[ \curlybraces{x} \]
\[ \curlybraces*{\frac{x}{3}} \]
\[ \curlybraces[\Bigg]{\frac{x}{3}} \]
```

$$\begin{aligned}\{\cdot\} \\ \{x\} \\ \left\{\frac{x}{3}\right\} \\ \left\{\frac{x}{3}\right\}\end{aligned}$$

<code>\magnitude[(size)]{&lt;quantity&gt;}</code>	(alias for double bars)
<code>\magnitude*[size]{&lt;quantity&gt;}</code>	(alias for double bars for fractions)
<code>\norm[(size)]{&lt;quantity&gt;}</code>	(alias for double bars)
<code>\norm*[size]{&lt;quantity&gt;}</code>	(alias for double bars for fractions)
<code>\absolutevalue[(size)]{&lt;quantity&gt;}</code>	(alias for single bars)
<code>\absolutevalue*[size]{&lt;quantity&gt;}</code>	(alias for single bars for fractions)

Semantic aliases. Use `\magnitude` or `\magnitude*` to typeset the magnitude of a vector.

```
\[ \magnitude{\vec{p}} \]
\[ \magnitude{\vec{p}} \]
\[ \magnitude*{\vec{p}_{\text{final}}} \]
\[ \magnitude*{\vec{p}_{\text{final}}} \]
```

$$\begin{aligned}\|p\| \\ \|\vec{p}\| \\ \|p_{\text{final}}\| \\ \|\vec{p}_{\text{final}}\|\end{aligned}$$

N 2021-04-06  
N 2021-04-06

\parallelto  
\perpendicularto

Commands for geometric relationships, mainly intended for subscripts.

```
\(\ \vec{F}_{\parallel}\{\parallelto} + \vec{F}_{\perp}\{\perpendicularto} \)
```

$F_{\parallel} + F_{\perp}$

## 5.2 Problems and Annotated Problem Solutions

N 2021-02-03

```
\begin{physicsproblem}{\langle title \rangle}
```

  \langle problem \rangle

(use this variant for vertical lists)

N 2021-02-03

```
\begin{physicsproblem*}{\langle title \rangle}
```

  \langle problem \rangle

(use this variant for in-line lists)

N 2021-02-03

```
\begin{parts}{\langle title \rangle}
```

  \langle problem \rangle

(provides problem parts)

```
\end{parts}
```

Provides an environment for stating physics problems. Each problem will begin on a new page. See the examples for how to handle single and multiple part problems.

N 2021-02-03

\problempart

Denotes a part of a problem within a `parts` environment.

```
\begin{physicsproblem}{Problem 1}
  This is a physics problem with no parts.
\end{physicsproblem}
```

### Problem 1

This is a physics problem with no parts.

```
\begin{physicsproblem}{Problem 2}
  This is a physics problem with multiple parts.
  The list is vertical.
  \begin{parts}
    \problempart This is the first part.
    \problempart This is the second part.
    \problempart This is the third part.
  \end{parts}
\end{physicsproblem}
```

## Problem 2

This is a physics problem with multiple parts. The list is vertical.

- (a) This is the first part.
- (b) This is the second part.
- (c) This is the third part.

```
\begin{physicsproblem*}{Problem 3}
  This is a physics problem with multiple parts.
  The list is in-line.
  \begin{parts}
    \problempart This is the first part.
    \problempart This is the second part.
    \problempart This is the third part.
  \end{parts}
\end{physicsproblem*}
```

## Problem 3

This is a physics problem with multiple parts. The list is in-line. (a) This is the first part. (b) This is the second part. (c) This is the third part.

**U** 2021-02-26

```
\begin{physicssolution}
  {solution steps}
\end{physicssolution}
\begin{physicssolution*}
  {solution steps}
\end{physicssolution*}
```

(use this variant for numbered steps)

**U** 2021-02-26

(use this variant for unnumbered steps)

This environment is only for mathematical solutions. The starred variant omits numbering of steps. See the examples.

<code>\begin{physicssolution} x &amp;= y + z \\ z &amp;= x - y \\ y &amp;= x - z \end{physicssolution}</code>	$x = y + z$ (1)
<code>\begin{physicssolution*} x &amp;= y + z \\ z &amp;= x - y \\ y &amp;= x - z \end{physicssolution*}</code>	$z = x - y$ (2)
	$y = x - z$ (3)
	$x = y + z$
	$z = x - y$
	$y = x - z$

U 2021-02-26

### \reason{<reason>}

Provides an annotation in a step-by-step solution. Keep reasons short and to the point. Wrap mathematical content in math mode.

<code>\begin{physicssolution} x &amp;= y + z \reason{This is a reason.} \\ z &amp;= x - y \reason{This is a reason too.} \\ y &amp;= x - z \reason{final answer} \end{physicssolution}</code>	$x = y + z$ This is a reason. (4)
<code>\begin{physicssolution*} x &amp;= y + z \reason{This is a reason.} \\ z &amp;= x - y \reason{This is a reason too.} \\ y &amp;= x - z \reason{final answer} \end{physicssolution*}</code>	$z = x - y$ This is a reason too. (5)
	$y = x - z$ final answer (6)
	$x = y + z$ This is a reason.
	$z = x - y$ This is a reason too.
	$y = x - z$ final answer

When writing solutions, remember that the `physicssolution`<sup>→P.56</sup> environment is *only* for mathematical content, not textual content or explanations.

```
\begin{physicsproblem}{Combined Problem and Solution}
This is an interesting physics problem.
\begin{physicssolution}
The solution goes here.
\end{physicssolution}
\end{physicsproblem}
```

```
\begin{physicsproblem}{Combined Multipart Problem with Solutions}
This is a physics problem with multiple parts.
\begin{parts}
\problempart This is the first part.
\begin{physicssolution}
The solution goes here.
\end{physicssolution}
\problempart This is the second part.
\begin{physicssolution}
The solution goes here.
\end{physicssolution}
\problempart This is the third part.
\begin{physicssolution}
The solution goes here.
\end{physicssolution}
\end{parts}
\end{physicsproblem}
```

N 2021-02-06

### \hilite[(color)]{<target>}[(shape)]

Hilites the desired target, which can be an entire mathematical expression or a part thereof. The default color is magenta and the default shape is a rectangle.

```
\begin{align*}
(\Delta s)^2 &= -(\Delta t)^2 + (\Delta x)^2 + (\Delta y)^2 + (\Delta z)^2 \\
&\quad + \\
(\Delta s)^2 &= \hilite{-(\Delta t)^2 + (\Delta x)^2}[rounded rectangle] + \\
&\quad (\Delta y)^2 + (\Delta z)^2 \\
(\Delta s)^2 &= \hilite{-(\Delta t)^2 + (\Delta x)^2}[rectangle] + \\
&\quad (\Delta y)^2 + (\Delta z)^2 \\
(\Delta s)^2 &= \hilite{-(\Delta t)^2 + (\Delta x)^2}[ellipse] + \\
&\quad (\Delta y)^2 + (\Delta z)^2 \\
(\Delta s)^2 &= \hilite{2}[circle] = \hilite[green]{-}[circle] \\
&\quad (\Delta t)^2 \hilite[cyan]{2}[circle] + \\
&\quad (\Delta x)^2 \hilite[orange]{2}[circle] + \\
&\quad (\Delta y)^2 \hilite[blue!50]{2}[circle] + \\
&\quad (\Delta z)^2 \hilite[violet!45]{2}[circle]
\end{align*}
```

$$\begin{aligned}
(\Delta s)^2 &= -(\Delta t)^2 + (\Delta x)^2 + (\Delta y)^2 + (\Delta z)^2 \\
(\Delta s)^2 &= -(\Delta t)^2 + (\Delta x)^2 + (\Delta y)^2 + (\Delta z)^2 \\
(\Delta s)^2 &= -(\Delta t)^2 + (\Delta x)^2 + (\Delta y)^2 + (\Delta z)^2 \\
(\Delta s)^2 &= -(\Delta t)^2 + (\Delta x)^2 + (\Delta y)^2 + (\Delta z)^2 \\
(\Delta s)^2 &= -(\Delta t)^2 + (\Delta x)^2 + (\Delta y)^2 + (\Delta z)^2
\end{aligned}$$

```
\begin{align*}
\Delta \vec{p} &= \vec{F}_{\text{net}} \Delta t \\
\text{\textcolor{orange}{hilite[orange]{\Delta \vec{p}}}} &\text{\textcolor{orange}{circle}} &= \vec{F}_{\text{net}} \Delta t \\
\Delta \vec{p} &= \text{\textcolor{yellow!50}{hilite[yellow!50]{\vec{F}_{\text{net}}}}} \Delta t \\
&\text{\textcolor{brown}{rounded rectangle}} \Delta t \\
\Delta \vec{p} &= \vec{F}_{\text{net}} \Delta t \\
&\text{\textcolor{brown}{rectangle}} \\
\Delta \vec{p} &= \text{\textcolor{cyan!50}{hilite[cyan!50]{\vec{F}_{\text{net}}}}} \Delta t \\
&\text{\textcolor{brown}{ellipse}} \\
\text{\textcolor{brown}{hilite{\Delta \vec{p}}}} &\text{\textcolor{brown}{rectangle}} &= \vec{F}_{\text{net}} \Delta t
\end{align*}
```

$$\Delta p = F_{\text{net}} \Delta t$$

Δp =  $F_{\text{net}} \Delta t$   
Δp =  $F_{\text{net}} \Delta t$

U 2021-09-18

`\image[<options>]{<caption>}{<label>}{<image>}`

Simplified interface for importing an image. The images are treated as floats, so they may not appear at the most logically intuitive place.

```
\image[scale=0.20]{example-image-1x1}
{Image shown 20 percent actual size.}{refffig1}
```



Figure 1: Image shown 20 percent actual size.

Figure `\ref{refffig1}` is nice.  
It's captioned `\nameref{refffig1}` and is on page `\pageref{refffig1}`.

Figure 1 is nice. It's captioned `Image shown 20 percent actual size` and is on page 59.

```
\image[scale=0.20,angle=45]{example-image-1x1}
{Image shown 20 percent actual size and rotated.}{refffig1}
```




Figure 2: Image shown 20 percent actual size and rotated.

`Figure \ref{reffig2} is nice.  
It's captioned \nameref{reffig2} and is on page \pageref{reffig2}.`

Figure 2 is nice. It's captioned [Image shown 20 percent actual size and rotated](#) and is on page [60](#).

### 5.3 Coordinate-Free and Index Notation

Beyond the current level of introductory physics, we need intelligent commands for typesetting vector and tensor symbols and components suitable for both coordinate-free and index notations.

`\colvec[(delimiter)]{(c1, ..., cn)}`  
`\rowvec[(delimiter)]{(c1, ..., cn)}`

Typesets column vectors and row vectors as numeric or symbolic components. There can be more than three components. The delimiter used in the list of components can be specified; the default is a comma. Units are not supported, so these are mainly for symbolic work.

<code>\[ \colvec{1,2,3} \]</code> <code>\[ \rowvec{1,2,3} \]</code> <code>\[ \colvec{x^0,x^1,x^2,x^3} \]</code> <code>\[ \rowvec{x_0,x_1,x_2,x_3} \]</code>	 <div style="display: flex; align-items: center; justify-content: space-around;"> <div style="text-align: center;"> <math display="block">\begin{pmatrix} 1 \\ 2 \\ 3 \end{pmatrix}</math> </div> <div style="text-align: center;"> <math display="block">(1 \quad 2 \quad 3)</math> </div> <div style="text-align: center;"> <math display="block">\begin{pmatrix} x^0 \\ x^1 \\ x^2 \\ x^3 \end{pmatrix}</math> </div> <div style="text-align: center;"> <math display="block">(x_0 \quad x_1 \quad x_2 \quad x_3)</math> </div> </div>
--	--

`\veccomp{<symbol>}`  
`\veccomp*{<symbol>}`  
`\tencomp{<symbol>}`  
`\tencomp*{<symbol>}`

(use this variant for coordinate-free vector notation)  
 (use this variant for index vector notation)  
 (use this variant for coordinate-free tensor notation)  
 (use this variant for index tensor notation)

Conforms to ISO 80000-2 notation.

```
\(\veccomp{r} \\
\(\veccomp{*}{r} \\
\(\tencomp{r} \\
\(\tencomp{*}{r}
```

**r**  
r  
**r**  
r

**\valence{<index>}{<index>}**  
**\valence\*{<index>}{<index>}**

Typesets tensor valence. The starred variant typesets it horizontally.

```
A vector is a \(\valence{1}{0} tensor. \\
A vector is a \(\valence*{1}{0} tensor.
```

A vector is a  $\begin{pmatrix} 1 \\ 0 \end{pmatrix}$  tensor.  
A vector is a  $(1, 0)$  tensor.

**\contraction{<slot,slot>}**  
**\contraction\*{<slot,slot>}**

Typesets tensor contraction in coordinate-free notation. There is no standard on this so we assert one here.

```
\(\contraction{1,2} \\
\(\contraction*{1,2}
```

$\mathbb{C}_{1,2}$   
 $C_{1,2}$

**\slot{<vector>}**  
**\slot\*{<vector>}**

An intelligent slot command for coordinate-free vector and tensor notation. The starred variants suppress the underscore.

```
\(\slot \\
\(\slot[\vec{a}] \\
\(\slot* \\
\(\slot*[\vec{a}]
```

$(\underline{\underline{\underline{}}})$   
 $(\underline{\underline{a}})$   
 $(\underline{\underline{\underline{\quad}}})$   
 $(\underline{\underline{a}})$

**\df**

Intelligent differential and exterior derivative operator.

```
\[
  \int x \, dx
]
[
  \int x \, \df{x}
]
[
  \int x \, \df*[x]
```

$\int x \, dx$   
 $\int x \, dx$   
 $\int x \, \mathbf{d}x$

## 5.4 Web VPython and VPython Program Listings

Web VPython<sup>3</sup> and VPython<sup>4</sup> are programming environments (both use Python) frequently used in introductory physics to introduce students for modeling physical systems. mandi makes including code listings very simple for students.

## 5.5 The `webvpythonblock` Environment

**U** 2022-01-27

```
\begin{webvpythonblock}[(options)](<link>){<caption>}  
  {Web VPython code}  
\end{webvpythonblock}
```

(now includes a QR code)

**N** 2022-01-27

```
\begin{webvpythonblock*}[(options)](<link>){<caption>}  
  {Web VPython code}  
\end{webvpythonblock*}
```

(use this variant to omit QR code)

Code placed here is nicely formatted and optionally linked to its source on [WebVPython.org](https://WebVPython.org), which must be in a public (not private) folder. Clicking anywhere in the code window will open the link in the default browser. A caption is mandatory, and a label is internally generated. The listing always begins on a new page. A URL shortening utility is recommended to keep the URL from getting unruly. For convenience, `https://` is automatically prepended to the URL and can thus be omitted. The `#` character in a URL should not cause problems. The default URL is that of the Web VPython home page.

---

<sup>3</sup>On November 9, 2021 GlowScript was renamed to Web VPython. The website was changed to <https://webvpython.org>.

<sup>4</sup><https://vpython.org>

```

\begin{webvpythonblock}(tinyurl.com/y3lnqyn3){A \texttt{Web VPython} Program With QR Code}
GlowScript 3.0 vpython

scene.width = 400
scene.height = 760
# constants and data
g = 9.8      # m/s^2
mball = 0.03 # kg
Lo = 0.26    # m
ks = 1.8     # N/m
deltat = 0.01 # s

# objects (origin is at ceiling)
ceiling = box(pos=vector(0,0,0), length=0.2, height=0.01,
               width=0.2)
ball = sphere(pos=vector(0,-0.3,0),radius=0.025,
              color=color.orange)
spring = helix(pos=ceiling.pos, axis=ball.pos-ceiling.pos,
               color=color.cyan,thickness=0.003,coils=40,
               radius=0.010)

# initial values
pball = mball * vector(0,0,0)      # kg m/s
Fgrav = mball * g * vector(0,-1,0) # N
t = 0

# improve the display
scene.autoscale = False           # turn off automatic camera zoom
scene.center = vector(0,-Lo,0) # move camera down
scene.waitfor('click')          # wait for a mouse click

# initial calculation loop
# calculation loop
while t < 10:
    rate(100)
    # we need the stretch
    s = mag(ball.pos) - Lo
    # we need the spring force
    Fspring = ks * s * -norm(spring.axis)
    Fnet = Fgrav + Fspring
    pball = pball + Fnet * deltat
    ball.pos = ball.pos + (pball / mball) * deltat
    spring.axis = ball.pos - ceiling.pos
    t = t + deltat
\end{webvpythonblock}

```



## Web VPython Program 1: A Web VPython Program With QR Code

```
1 GlowScript 3.0 vpython
2
3 scene.width = 400
4 scene.height = 760
5 # constants and data
6 g = 9.8          # m/s^2
7 mball = 0.03    # kg
8 Lo = 0.26        # m
9 ks = 1.8         # N/m
10 deltat = 0.01   # s
11
12 # objects (origin is at ceiling)
13 ceiling = box(pos=vector(0,0,0), length=0.2, height=0.01,
14                 width=0.2)
15 ball = sphere(pos=vector(0,-0.3,0), radius=0.025,
16                 color=color.orange)
17 spring = helix(pos=ceiling.pos, axis=ball.pos-ceiling.pos,
18                  color=color.cyan, thickness=0.003, coils=40,
19                  radius=0.010)
20
21 # initial values
22 pball = mball * vector(0,0,0)      # kg m/s
23 Fgrav = mball * g * vector(0,-1,0) # N
24 t = 0
25
26 # improve the display
27 scene.autoscale = False           # turn off automatic camera zoom
28 scene.center = vector(0,-Lo,0)     # move camera down
29 scene.waitfor('click')            # wait for a mouse click
30
31 # initial calculation loop
32 # calculation loop
33 while t < 10:
34     rate(100)
35     # we need the stretch
36     s = mag(ball.pos) - Lo
37     # we need the spring force
38     Fspring = ks * s * -norm(spring.axis)
39     Fnet = Fgrav + Fspring
40     pball = pball + Fnet * deltat
41     ball.pos = ball.pos + (pball / mball) * deltat
42     spring.axis = ball.pos - ceiling.pos
43     t = t + deltat
```

Here is how one would reference this program elsewhere. Notice the references are numbered sequentially within the document.

\WebVPython\ program \ref{gs:1} is nice.  
It's called \nameref{gs:1} and is on page \pageref{gs:1}.

Web VPython program 1 is nice. It's called [A Web VPython Program With QR Code](#) and is on page 64.

```

\begin{webvpythonblock*}(tinyurl.com/y3lnqyn3){A \texttt{Web VPython} Program Without QR Code}
GlowScript 3.0 vpython

scene.width = 400
scene.height = 760
# constants and data
g = 9.8      # m/s^2
mball = 0.03 # kg
Lo = 0.26    # m
ks = 1.8     # N/m
deltat = 0.01 # s

# objects (origin is at ceiling)
ceiling = box(pos=vector(0,0,0), length=0.2, height=0.01,
               width=0.2)
ball = sphere(pos=vector(0,-0.3,0),radius=0.025,
              color=color.orange)
spring = helix(pos=ceiling.pos, axis=ball.pos-ceiling.pos,
                color=color.cyan,thickness=0.003,coils=40,
                radius=0.010)

# initial values
pball = mball * vector(0,0,0)      # kg m/s
Fgrav = mball * g * vector(0,-1,0) # N
t = 0

# improve the display
scene.autoscale = False           # turn off automatic camera zoom
scene.center = vector(0,-Lo,0) # move camera down
scene.waitfor('click')           # wait for a mouse click

# initial calculation loop
# calculation loop
while t < 10:
    rate(100)
    # we need the stretch
    s = mag(ball.pos) - Lo
    # we need the spring force
    Fspring = ks * s * -norm(spring.axis)
    Fnet = Fgrav + Fspring
    pball = pball + Fnet * deltat
    ball.pos = ball.pos + (pball / mball) * deltat
    spring.axis = ball.pos - ceiling.pos
    t = t + deltat
\end{webvpythonblock*}

```

## Web VPython Program 2: A Web VPython Program Without QR Code

```
1 GlowScript 3.0 vpython
2
3 scene.width = 400
4 scene.height = 760
5 # constants and data
6 g = 9.8      # m/s^2
7 mball = 0.03 # kg
8 Lo = 0.26    # m
9 ks = 1.8     # N/m
10 deltat = 0.01 # s
11
12 # objects (origin is at ceiling)
13 ceiling = box(pos=vector(0,0,0), length=0.2, height=0.01,
14                 width=0.2)
15 ball = sphere(pos=vector(0,-0.3,0), radius=0.025,
16                 color=color.orange)
17 spring = helix(pos=ceiling.pos, axis=ball.pos-ceiling.pos,
18                  color=color.cyan, thickness=0.003, coils=40,
19                  radius=0.010)
20
21 # initial values
22 pball = mball * vector(0,0,0)      # kg m/s
23 Fgrav = mball * g * vector(0,-1,0) # N
24 t = 0
25
26 # improve the display
27 scene.autoscale = False           # turn off automatic camera zoom
28 scene.center = vector(0,-Lo,0) # move camera down
29 scene.waitfor('click')          # wait for a mouse click
30
31 # initial calculation loop
32 # calculation loop
33 while t < 10:
34     rate(100)
35     # we need the stretch
36     s = mag(ball.pos) - Lo
37     # we need the spring force
38     Fspring = ks * s * -norm(spring.axis)
39     Fnet = Fgrav + Fspring
40     pball = pball + Fnet * deltat
41     ball.pos = ball.pos + (pball / mball) * deltat
42     spring.axis = ball.pos - ceiling.pos
43     t = t + deltat
```

\WebVPython\ program \ref{gs:2} is nice.  
It's called \nameref{gs:2} and is on page \pageref{gs:2}.

Web VPython program 2 is nice. It's called [A Web VPython Program Without QR Code](#) and is on page [67](#).

## 5.6 The vpythonfile Command

U 2022-01-27

\vpythonfile[*options*] (*link*) {*file*} {*caption*}

Command to load and typeset a VPython program, read from local file {*file*}. Clicking anywhere in the code window can optionally open a link, passed as an option, in the default browser. A caption is mandatory, and a label is internally generated. The listing always begins on a new page. A URL shortening utility is recommended to keep the URL from getting unruly. For convenience, <https://> is automatically prepended to the URL and can thus be omitted. The default URL is that of the VPython home page.

```
\vpythonfile{vdemo.py}{A \VPython\ Program}
```

## VPython Program 1: A VPython Program

```
1  from vpython import *
2
3  scene.width = 400
4  scene.height = 760
5  # constants and data
6  g = 9.8      # m/s^2
7  mball = 0.03 # kg
8  Lo = 0.26    # m
9  ks = 1.8     # N/m
10 deltat = 0.01 # s
11
12 # objects (origin is at ceiling)
13 ceiling = box(pos=vector(0,0,0), length=0.2, height=0.01,
14                 width=0.2)
15 ball = sphere(pos=vector(0,-0.3,0), radius=0.025,
16                 color=color.orange)
17 spring = helix(pos=ceiling.pos, axis=ball.pos-ceiling.pos,
18                  color=color.cyan, thickness=0.003, coils=40,
19                  radius=0.010)
20
21 # initial values
22 pball = mball * vector(0,0,0)      # kg m/s
23 Fgrav = mball * g * vector(0,-1,0) # N
24 t = 0
25
26 # improve the display
27 scene.autoscale = False           # turn off automatic camera zoom
28 scene.center = vector(0,-Lo,0)    # move camera down
29 scene.waitfor('click')           # wait for a mouse click
30
31 # initial calculation loop
32 # calculation loop
33 while t < 10:
34     rate(100)
35     # we need the stretch
36     s = mag(ball.pos) - Lo
37     # we need the spring force
38     Fspring = ks * s * -norm(spring.axis)
39     Fnet = Fgrav + Fspring
40     pball = pball + Fnet * deltat
41     ball.pos = ball.pos + (pball / mball) * deltat
42     spring.axis = ball.pos - ceiling.pos
43     t = t + deltat
```

\VPython\ program \ref{vp:1} is nice.  
It's called \nameref{vp:1} and is on page \pageref{vp:1}.

VPython program 1 is nice. It's called [A VPython Program](#) and is on page [69](#).

## 5.7 The `webvpythoninline` and `vpythoninline` Commands

U 2021-02-26

`\webvpythoninline{< Web VPython code>}`

U 2021-02-26

`\vpythoninline{< VPython code>}`

Typesets a small, in-line snippet of code. The snippet should be less than one line long.

`\WebVPython\ programs begin with \webvpythoninline{GlowScript 3.0 VPython}`  
`and \VPython\ programs begin with \vpythoninline{from vpython import *}.`

Web VPython programs begin with `GlowScript 3.0 VPython` and VPython programs begin with  
`from vpython import *`.

## 5.8 **mandistudent** Source Code

Define the package version and date for global use, exploiting the fact that in a .sty file there is now no need for `\makeatletter` and `\makeatother`. This simplifies defining internal commands, with @ in the name, that are not for the user to know about.

```

1 \def\mandistudent@version{3.1.0}
2 \def\mandistudent@date{2022-01-27}
3 \NeedsTeXFormat{LaTeXe}[2020-02-02]
4 \DeclareRelease{v3.1.0}{2022-01-27}{mandistudent.sty}
5 \DeclareCurrentRelease{v\mandistudent@version}{\mandistudent@date}
6 \ProvidesPackage{mandistudent}
7   [\mandistudent@date\space v\mandistudent@version\space Macros for introductory physics]
```

Define a convenient package version command.

```
8 \newcommand*\mandistudentversion{v\mandistudent@version\space dated \mandistudent@date}
```

Load third party packages, documenting why each one is needed.

```

9 \RequirePackage{amsmath}           % AMS goodness (don't load amssymb or amsfonts)
10 \RequirePackage[inline]{enumitem}  % needed for physicsproblem environment
11 \RequirePackage{eso-pic}          % needed for \hilite
12 \RequirePackage[g]{esvect}        % needed for nice vector arrow, style g
13 \RequirePackage{pgfopts}          % needed for key-value interface
14 \RequirePackage{iftex}            % needed for requiring LuaLaTeX
15 \RequirePackage{makebox}          % needed for consistent \dirvect; \makebox
16 \RequirePackage{mandi}            %
17 \RequirePackage{mathtools}         % needed for paired delimiters; extends amsmath
18 \RequirePackage{nicematrix}        % needed for column and row vectors
19 \RequirePackage{qrcode}           % needed for QR codes in webpythonblock
20 \qrset{height=1.5cm}             % set default size of QR code
21 \RequirePackage[most]{tcolorbox}  % needed for program listings
22 \RequirePackage{tensor}           % needed for index notation
23 \RequirePackage{tikz}              % needed for \hilite
24 \usetikzlibrary{shapes,fit,tikzmark} % needed for \hilite
25 \RequirePackage{unicode-math}     % needed for Unicode support

26 \IfFormatAtLeastTF {2020-10-01}  % load xparse if necessary
27   {}%
28   {\RequirePackage{xparse}}%
29 \RequirePackage{hyperref}          % load last
30 \RequireLuaTeX                  % require this engine
```

Set up the fonts to be consistent with ISO 80000-2 notation. The `unicode-math` package loads the `fontspec` and `xparse` packages. Note that `xparse` is now part of the L<sup>A</sup>T<sub>E</sub>X 2 <sub>$\epsilon$</sub>  kernel. Because `unicode-math` is required, all documents using `mandi` must be compiled with an engine that supports Unicode. We recommend LuaL<sup>A</sup>T<sub>E</sub>X.

```
31 \unimathsetup{math-style=ISO}
32 \unimathsetup{warnings-off={mathtools-colon,mathtools-overbracket}}
```

Use normal math letters from Latin Modern Math for familiarity with textbooks.

```
33 \setmathfont[Scale=MatchLowercase]
34   {Latin Modern Math}  % default math font; better J
```

Borrow from TeX Gyre DejaVu Math for vectors and tensors to get single-storey g.

```
35 \setmathfont[Scale=MatchLowercase,range={sfit/{latin},bfsfit/{latin}}]
36   {TeX Gyre DejaVu Math} % single-storey lowercase g
```

Borrow from TeX Gyre DejaVu Math to get single-storey g.

```
37 \setmathfont[Scale=MatchLowercase,range={sfup/{latin},bfsfup/{latin}}]
38   {TeX Gyre DejaVu Math} % single-storey lowercase g
```

Borrow `\mathscr` and `\mathbf{\mathscr}` from XITS Math.

See <https://tex.stackexchange.com/a/120073/218142>.

```
39 \setmathfont[Scale=MatchLowercase,range={\mathscr,\mathbf{\mathscr}}]{XITS Math}
```

Get original and bold `\mathcal` fonts.

See <https://tex.stackexchange.com/a/21742/218142>.

```
40 \setmathfont[Scale=MatchLowercase,range={\mathcal,\mathbf{\mathcal}},StylisticSet=1]{XITS Math}
```

Borrow Greek sfup and sfit letters from STIX Two Math. Since this isn't officially supported in `unicode-math` we have to manually set this up.

```
41 \setmathfont[Scale=MatchLowercase,range={"E17C-"E1F6}]{STIX Two Math}
```

```
42 \newfontfamily{\symsfgreek}{STIX Two Math}
```

```
43 % I don't understand why \text{...} is necessary.
```

```
44 \newcommand{\symsfupalpha}{\text{\symsfgreek{^\wedge e196}}}
45 \newcommand{\symsfupbeta}{\text{\symsfgreek{^\wedge e197}}}
46 \newcommand{\symsfupgamma}{\text{\symsfgreek{^\wedge e198}}}
47 \newcommand{\symsfupdelta}{\text{\symsfgreek{^\wedge e199}}}
48 \newcommand{\symsfupepsilon}{\text{\symsfgreek{^\wedge e1af}}}
49 \newcommand{\symsfupvarepsilon}{\text{\symsfgreek{^\wedge e19a}}}
50 \newcommand{\symsfupzeta}{\text{\symsfgreek{^\wedge e19b}}}
51 \newcommand{\symsfupeta}{\text{\symsfgreek{^\wedge e19c}}}
52 \newcommand{\symsfuptheta}{\text{\symsfgreek{^\wedge e19d}}}
53 \newcommand{\symsfupvartheta}{\text{\symsfgreek{^\wedge e1b0}}}
54 \newcommand{\symsfupiota}{\text{\symsfgreek{^\wedge e19e}}}
55 \newcommand{\symsfupkappa}{\text{\symsfgreek{^\wedge e19f}}}
56 \newcommand{\symsfuplambda}{\text{\symsfgreek{^\wedge e1a0}}}
57 \newcommand{\symsfupmu}{\text{\symsfgreek{^\wedge e1a1}}}
58 \newcommand{\symsfupnu}{\text{\symsfgreek{^\wedge e1a2}}}
59 \newcommand{\symsfupxi}{\text{\symsfgreek{^\wedge e1a3}}}
60 \newcommand{\symsfupomicron}{\text{\symsfgreek{^\wedge e1a4}}}
61 \newcommand{\symsfuppi}{\text{\symsfgreek{^\wedge e1a5}}}
62 \newcommand{\symsfupvarpi}{\text{\symsfgreek{^\wedge e1b3}}}
63 \newcommand{\symsfuprho}{\text{\symsfgreek{^\wedge e1a6}}}
64 \newcommand{\symsfupvarrho}{\text{\symsfgreek{^\wedge e1b2}}}
65 \newcommand{\symsfupsigma}{\text{\symsfgreek{^\wedge e1a8}}}
66 \newcommand{\symsfupvarsigma}{\text{\symsfgreek{^\wedge e1a7}}}
67 \newcommand{\symsfuptau}{\text{\symsfgreek{^\wedge e1a9}}}
68 \newcommand{\symsfupupsilon}{\text{\symsfgreek{^\wedge e1aa}}}
69 \newcommand{\symsfupphi}{\text{\symsfgreek{^\wedge e1b1}}}
70 \newcommand{\symsfupvarphi}{\text{\symsfgreek{^\wedge e1ab}}}
71 \newcommand{\symsfupchi}{\text{\symsfgreek{^\wedge e1ac}}}
72 \newcommand{\symsfuppsi}{\text{\symsfgreek{^\wedge e1ad}}}
73 \newcommand{\symsfupomega}{\text{\symsfgreek{^\wedge e1ae}}}
74 \newcommand{\symsfupDelta}{\text{\symsfgreek{^\wedge e180}}}
75 \newcommand{\symsfupGamma}{\text{\symsfgreek{^\wedge e17f}}}
76 \newcommand{\symsfupTheta}{\text{\symsfgreek{^\wedge e18e}}}
77 \newcommand{\symsfupLambda}{\text{\symsfgreek{^\wedge e187}}}
78 \newcommand{\symsfupXi}{\text{\symsfgreek{^\wedge e18a}}}
79 \newcommand{\symsfupPi}{\text{\symsfgreek{^\wedge e18c}}}
80 \newcommand{\symsfupSigma}{\text{\symsfgreek{^\wedge e18f}}}
81 \newcommand{\symsfupUpsilon}{\text{\symsfgreek{^\wedge e191}}}
82 \newcommand{\symsfupPhi}{\text{\symsfgreek{^\wedge e192}}}
83 \newcommand{\symsfupPsi}{\text{\symsfgreek{^\wedge e194}}}
84 \newcommand{\symsfupOmega}{\text{\symsfgreek{^\wedge e195}}}
85 \newcommand{\symsfitalpha}{\text{\symsfgreek{^\wedge e1d8}}}
86 \newcommand{\symsfitbeta}{\text{\symsfgreek{^\wedge e1d9}}}
87 \newcommand{\symsfitgamma}{\text{\symsfgreek{^\wedge e1da}}}
88 \newcommand{\symsfitdelta}{\text{\symsfgreek{^\wedge e1db}}}
```

```

89 \newcommand{\symsfitepsilon} {\text{\symsfgreek{e1f1}}}
90 \newcommand{\symsfitvarepsilon} {\text{\symsfgreek{e1dc}}}
91 \newcommand{\symsfitzeta} {\text{\symsfgreek{e1dd}}}
92 \newcommand{\symsfiteta} {\text{\symsfgreek{e1de}}}
93 \newcommand{\symsfittheta} {\text{\symsfgreek{e1df}}}
94 \newcommand{\symsfitvartheta} {\text{\symsfgreek{e1f2}}}
95 \newcommand{\symsfitiota} {\text{\symsfgreek{e1e0}}}
96 \newcommand{\symsfitkappa} {\text{\symsfgreek{e1e1}}}
97 \newcommand{\symsfitlambda} {\text{\symsfgreek{e1e2}}}
98 \newcommand{\symsfitmu} {\text{\symsfgreek{e1e3}}}
99 \newcommand{\symsfitnu} {\text{\symsfgreek{e1e4}}}
100 \newcommand{\symsfitxi} {\text{\symsfgreek{e1e5}}}
101 \newcommand{\symsfitomicron} {\text{\symsfgreek{e1e6}}}
102 \newcommand{\symsfitpi} {\text{\symsfgreek{e1e7}}}
103 \newcommand{\symsfitvarpi} {\text{\symsfgreek{e1f5}}}
104 \newcommand{\symsfitrho} {\text{\symsfgreek{e1e8}}}
105 \newcommand{\symsfitvarrho} {\text{\symsfgreek{e1f4}}}
106 \newcommand{\symsfitsigma} {\text{\symsfgreek{e1ea}}}
107 \newcommand{\symsfitvarsigma} {\text{\symsfgreek{e1e9}}}
108 \newcommand{\symsfittau} {\text{\symsfgreek{e1eb}}}
109 \newcommand{\symsfitupsilon} {\text{\symsfgreek{e1ec}}}
110 \newcommand{\symsfitphi} {\text{\symsfgreek{e1f3}}}
111 \newcommand{\symsfitvarphi} {\text{\symsfgreek{e1ed}}}
112 \newcommand{\symsfitchi} {\text{\symsfgreek{e1ee}}}
113 \newcommand{\symsfitpsi} {\text{\symsfgreek{e1ef}}}
114 \newcommand{\symsfitomega} {\text{\symsfgreek{e1f0}}}
115 \newcommand{\symsfitDelta} {\text{\symsfgreek{e1c2}}}
116 \newcommand{\symsfitGamma} {\text{\symsfgreek{e1c1}}}
117 \newcommand{\symsfitTheta} {\text{\symsfgreek{e1d0}}}
118 \newcommand{\symsfitLambda} {\text{\symsfgreek{e1c9}}}
119 \newcommand{\symsfitXi} {\text{\symsfgreek{e1cc}}}
120 \newcommand{\symsfitPi} {\text{\symsfgreek{e1ce}}}
121 \newcommand{\symsfitSigma} {\text{\symsfgreek{e1d1}}}
122 \newcommand{\symsfitUpsilon} {\text{\symsfgreek{e1d3}}}
123 \newcommand{\symsfitPhi} {\text{\symsfgreek{e1d4}}}
124 \newcommand{\symsfitPsi} {\text{\symsfgreek{e1d6}}}
125 \newcommand{\symsfitOmega} {\text{\symsfgreek{e1d7}}}

```

Tweak the `esvect` package fonts to get the correct font size.

See <https://tex.stackexchange.com/a/566676>.

```

126 \DeclareFontFamily{U}{esvect}{}%
127 \DeclareFontShape{U}{esvect}{m}{n}{%
128   <-5.5> vect5
129   <5.5-6.5> vect6
130   <6.5-7.5> vect7
131   <7.5-8.5> vect8
132   <8.5-9.5> vect9
133   <9.5-> vect10
134 }{}%

```

Write a banner to the console showing the options in use.

```

135 \typeout{}%
136 \typeout{mandistudent: You are using mandistudent \mandistudentversion.}%
137 \typeout{mandistudent: This package requires LuaTeX.}%
138 \typeout{mandistudent: This package changes the default math font(s).}%
139 \typeout{mandistudent: This package redefines the \protect\vec\space command.}%
140 \typeout{}%

```

A better, intelligent coordinate-free  $\text{\vec}^{\text{P.51}}$  command. Note the use of the  $\text{\vec}_{\text{~}}$  type of optional argument. This accounts for much of the flexibility and power of this command. Also note the use of the TeX primitives `\sb{}` and `\sp{}`.

Why doesn't it work when I put spaces around #3 or #4? Because outside of `\ExplSyntaxOn... \ExplSyntaxOff`, the `_` character has a different catcode and is treated as a mathematical entity.

See <https://tex.stackexchange.com/q/554706/218142>.

See also <https://tex.stackexchange.com/a/531037/218142>.

```

141 \RenewDocumentCommand{\vec}{ s m e{_} }%
142   {%
143     % Note the \, used to make superscript look better.
144     \IfBooleanTF{#1}%
145       {%
146         \vv{#2}%      % * gives an arrow
147         % Use \sp{} primitive for superscript.
148         % Adjust superscript for the arrow.
149         \IfValueT{#4}%
150           {\sp{\, #4\vphantom{\smash[t]{\big|}}}}}
151       }%
152     {%
153       \symbfit{#2} % no * gives us bold
154       % Use \sp{} primitive for superscript.
155       % No superscript adjustment needed.
156       \IfValueT{#4}%
157         {\sp{#4\vphantom{\smash[t]{\big|}}}}
158     }%
159     % Use \sb{} primitive for subscript.
160     \IfValueT{#3}%
161       {\sb{#3\vphantom{\smash[b]{|}}}}
162   }%

```

A command for the direction of a vector. We use a slight tweak to get uniform hats that requires the `makebox` package.

See <https://tex.stackexchange.com/a/391204/218142>.

```

163 \NewDocumentCommand{\dirvec}{ s m e{_} }%
164   {%
165     \widehat{%
166       \makebox*{(\w\!)}%
167     }%
168     \ensuremath{%
169       \IfBooleanTF {#1}%
170         {%
171           #2%
172         }%
173       }%
174     \symbfit{#2}%
175   }%
176   }%
177   }%
178 }%
179 }%
180 \IfValueT{#3}%
181   {\sb{#3\vphantom{\smash[b]{|}}}}%
182 \IfValueT{#4}%
183   {\sp{\, #4\vphantom{\smash[t]{\big|}}}}%
184 }%

```

The zero vector.

```

185 \NewDocumentCommand{\zerovec}{ s }%
186   {%
187     \IfBooleanTF {#1}%
188       {\vv{0}}%
189       {\symbup{0}}%
190   }%

```

Notation for column and row vectors.

See <https://tex.stackexchange.com/a/39054/218142>.

```
191 \ExplSyntaxOn
192 \NewDocumentCommand{\colvec}{ O{,} m }
193 {
194     \__mandi_vectormain:n { p } { \\\ } { #1 } { #2 }
195 }
196 \NewDocumentCommand{\rowvec}{ O{,} m }
197 {
198     \__mandi_vectormain:n { p } { & } { #1 } { #2 }
199 }
200 \seq_new:N \l__mandi_vectorarg_seq
201 \cs_new_protected:Npn \__mandi_vectormain:n { p } { #1#2#3#4 }
202 {
203     \seq_set_split:Nnn \l__mandi_vectorarg_seq { #3 } { #4 }
204     \begin{#1NiceMatrix}[r]
205         \seq_use:Nnnn \l__mandi_vectorarg_seq { #2 } { #2 } { #2 }
206     \end{#1NiceMatrix}
207 }
208 \ExplSyntaxOff
```

Students always need this symbol.

```
209 \NewDocumentCommand{\changein}{\Delta}
```

Intelligent delimiters provided via the `mathtools` package. Use the starred variants for fractions. You can supply optional sizes. Note that default placeholders are used when the argument is empty.

```
210 \DeclarePairedDelimiterX{\doublebars}[1]{\lVert}{\rVert}{\ifblank{#1}{\cdot}{\cdotp\cdotp\cdotp}}
211 \DeclarePairedDelimiterX{\singlebars}[1]{\lvert}{\rvert}{\ifblank{#1}{\cdot}{\cdotp\cdotp\cdotp}}
212 \DeclarePairedDelimiterX{\anglebrackets}[1]{\langle}{\rangle}{\ifblank{#1}{\cdot}{\cdotp\cdotp\cdotp}}
213 \DeclarePairedDelimiterX{\parentheses}[1]{(}{)}{\ifblank{#1}{\cdot}{\cdotp\cdotp\cdotp}}
214 \DeclarePairedDelimiterX{\squarebrackets}[1]{[}{]}{\ifblank{#1}{\cdot}{\cdotp\cdotp\cdotp}}
215 \DeclarePairedDelimiterX{\curlybraces}[1]{\{}{\}}{\ifblank{#1}{\cdot}{\cdotp\cdotp\cdotp}}
```

Some semantic aliases. Because of the way `\vecP.51` and `\dirvecP.52` are defined, I reluctantly decided not to implement a `\magvec` command. It would require accounting for too many options. So `\magnitudeP.54` is the new solution.

```
216 \NewDocumentCommand{\magnitude}{\doublebars}
217 \NewDocumentCommand{\norm}{\doublebars}
218 \NewDocumentCommand{\absolutevalue}{\singlebars}
```

Commands for two important geometric relationships. These are meant mainly to be subscripts.

```
219 \NewDocumentCommand{\parallelto}{\doublebars}{%
220     \mkern3mu\vphantom{\perp}\vrule depth 0pt\mkern2mu\vrule depth 0pt\mkern3mu%
221 }
222 \NewDocumentCommand{\perpendicularto}{\perp}
```

An environment for problem statements. The starred variant gives in-line lists.

```
224 \NewDocumentEnvironment{physicsproblem}{ m }{%
225     \newpage%
226     \section*{#1}%
227     \newlist{parts}{enumerate}{2}%
228     \setlist[parts]{label=\bfseries(\alph*)}%
229 }
230 \NewDocumentEnvironment{physicsproblem*}{ m }{%
231     \newpage%
232 }
```

```

235 \section*{[#1]%
236 \newlist{parts}{enumerate*}{2}%
237 \setlist[parts]{label=\bfseries(\alph*)}%
238 }%
239 {}%
240 \NewDocumentCommand{\problempart}{\item}%

An environment for problem solutions.

241 \NewDocumentEnvironment{physicssolution}{ +b }{%
242   f%
243   % Make equation numbering consecutive through the document.
244   \begin{align}%
245     #1%
246   \end{align}%
247 }%
248 {}%
249 \NewDocumentEnvironment{physicssolution*}{ +b }{%
250   f%
251   % Make equation numbering consecutive through the document.
252   \begin{align*}%
253     #1%
254   \end{align*}%
255 }%
256 {}%

```

See <https://tex.stackexchange.com/q/570223/218142>.

```

257 \NewDocumentCommand{\reason}{ 0{4cm} m }{%
258   f%
259   &&\begin{minipage}{#1}\raggedright\small #2\end{minipage}%
260 }%

```

Command for highlighting parts of, or entire, mathematical expressions.

See <https://texample.net/tikz/examples/beamer-arrows/>.

See also <https://tex.stackexchange.com/a/406084/218142>.

See also <https://tex.stackexchange.com/a/570858/218142>.

See also <https://tex.stackexchange.com/a/570789/218142>.

See also <https://tex.stackexchange.com/a/79659/218142>.

See also <https://tex.stackexchange.com/q/375032/218142>.

See also <https://tex.stackexchange.com/a/571744/218142>

```

261 \newcounter{tikzhighlightnode}%
262 \NewDocumentCommand{\hilite}{ 0{magenta!60} m 0{rectangle} }{%
263   f%
264   \stepcounter{tikzhighlightnode}%
265   \tikzmarknode{highlighted-node-\number\value{tikzhighlightnode}}{#2}%
266   \edef\temp{%
267     \noexpand\AddToShipoutPictureBG{%
268       \noexpand\begin{tikzpicture}[overlay,remember picture]%
269       \noexpand\iftikzmarkoncurrentpage{highlighted-node-\number\value{tikzhighlightnode}}{%
270         \noexpand\node[inner sep=1.0pt,fill=#1,#3,fit=(highlighted-node-\number\value{tikzhighlightnode})]{};%
271       \noexpand\fi%
272       \noexpand\end{tikzpicture}%
273     }%
274   }%
275   \temp%
276 }%

```

A simplified command for importing images.

See <https://tex.stackexchange.com/a/614478/218142>.

```

277 \NewDocumentCommand{\image}{ O{scale=1} m m m }%
278   {%
279     \par
280     \begin{figure}[ht!]
281       \centering%
282       \includegraphics[#1]{#2}%
283       \caption{#3}%
284       \label{#4}%
285     \end{figure}%
286     \par
287   }%

```

Intelligent commands for typesetting vector and tensor symbols and components suitable for use with both coordinate-free and index notations. Use starred form for index notation, unstarred form for coordinate-free.

```

288 \NewDocumentCommand{\veccomp}{ s m }%
289   {%
290     % Consider renaming this to \vectorsym.
291     \IfBooleanTF{#1}%
292       {%
293         \symnormal{#2}%
294       }%
295       {%
296         \symbfit{#2}%
297       }%
298   }%
299 \NewDocumentCommand{\tencomp}{ s m }%
300   {%
301     % Consider renaming this to \tensororsym.
302     \IfBooleanTF{#1}%
303       {%
304         \symsfit{#2}%
305       }%
306       {%
307         \symbfsfit{#2}%
308       }%
309   }%

```

Command to typeset tensor valence.

```

310 \NewDocumentCommand{\valence}{ s m m }%
311   {%
312     \IfBooleanTF{#1}%
313       {%
314         (#2,#3)%
315       }%
316       {%
317         \binom{#2}{#3}%
318       }%
319   }%

```

Intelligent notation for contraction on pairs of slots.

```

320 \NewDocumentCommand{\contraction}{ s m }%
321   {%
322     \IfBooleanTF{#1}%
323       {%
324         \mathsf{C}%
325       }%
326       {%
327         \mathbb{C}%
328       }%

```

```

329     _{#2}
330 }%
Intelligent slot command for coordinate-free tensor notation.

331 \NewDocumentCommand{\slot}{ s d[] }%
332 {%
333     % d[] must be used because of the way consecutive optional
334     % arguments are handled. See xparse docs for details.
335     \IfBooleanTF{#1}%
336     {%
337         \IfValueTF{#2}%
338             {%
339                 \smash{\makebox[1.5em]{\ensuremath{#2}}}%
340             }%
341             {%
342                 \smash{\makebox[1.5em]{\ensuremath{}}}%
343             }%
344     }%
345     {%
346         \IfValueTF{#2}%
347             {%
348                 \underline{\smash{\makebox[1.5em]{\ensuremath{#2}}}}%
349             }%
350             {%
351                 \underline{\smash{\makebox[1.5em]{\ensuremath{}}}}%
352             }%
353     }%
354 }%

```

Intelligent differential (exterior derivative) operator.

```

355 \NewDocumentCommand{\df}{ s }%
356 {%
357     \mathop{} \! %
358     \IfBooleanTF{#1}%
359     {%
360         \symbfsup{d}%
361     }%
362     {%
363         \symsup{d}%
364     }%
365 }%

```

Here is a clever way to color digits in program listings thanks to Ulrike Fischer.

See <https://tex.stackexchange.com/a/570717/218142>.

```

366 \directlua{%
367   luactfload.add_colorscheme("colordigits",
368     {[ "8000FF"] = {"one", "two", "three", "four", "five", "six", "seven", "eight", "nine", "zero"} })
369 }%
370 \newfontfamily\colordigits{DejaVuSansMono}[RawFeature={color=colordigits}]

```

Set up a color scheme and a new code environment for listings. The new colors are more restful on the eye. All listing commands now use `tcolorbox`.

See <https://tex.stackexchange.com/a/529421/218142>.

```

371 \newfontfamily{\gsfontfamily}{DejaVuSansMono}    % new font for listings
372 \definecolor{gsbggray}{rgb}{0.90,0.90,0.90} % background gray
373 \definecolor{gsgray}{rgb}{0.30,0.30,0.30} % gray
374 \definecolor{gsgreen}{rgb}{0.00,0.60,0.00} % green
375 \definecolor{gsorange}{rgb}{0.80,0.45,0.12} % orange
376 \definecolor{gspeach}{rgb}{1.00,0.90,0.71} % peach

```

```

377 \definecolor{gspearl}      {rgb}{0.94,0.92,0.84} % pearl
378 \definecolor{gsplum}        {rgb}{0.74,0.46,0.70} % plum
379 \lstdefinestyle{vpython}%
380   {%
381     backgroundcolor=\color{gsbggray},%           % style for listings
382     basicstyle=\color{gsblack}\footnotesize,%       % background color
383     breakatwhitespace=true%,                   % default style
384     breaklines=true,%                         % break at whitespace
385     captionpos=b,%                           % break long lines
386     classoffset=1,%                          % position caption
387     commentstyle=\color{gsgray},%                % STILL DON'T UNDERSTAND THIS
388     deletekeywords={print},%                  % font for comments
389     emph={self,cls,@classmethod,@property},%    % delete keywords from the given language
390     emphstyle=\color{gsorange}\itshape,%          % words to emphasize
391     escapeinside={(*@}{@*)},%                 % font for emphasis
392     frame=tb,%                                % add LaTeX within your code
393     framerule=2.0pt,%                         % frame style
394     framexleftmargin=5pt,%                     % frame thickness
395     identifierstyle=\sffamily,%                % extra frame left margin
396     keywordstyle=\sf\color{gsplum},%             % style for identifiers
397     language=Python,%                         % color for keywords
398     linewidth=\linewidth,%                   % select language
399     morekeywords=%                           % width of listings
400     __future__,abs,acos,align,ambient,angle,append,append_to_caption,% % VPython/Web VPython specific keywords
401     append_to_title,arange,arrow,asin,astuple,atan,atan2,attach_arrow,% %
402     attach_trail,autoscale,axis,background,billboard,bind,black,blue,border,% %
403     bounding_box,box,bumpaxis,bumpmap,bumpmaps,camera,canvas,caption,capture,% %
404     ceil,center,clear,clear_trail,click,clone,CoffeeScript,coils,color,combin,% %
405     comp,compound,cone,convex,cos,cross,curve,cyan,cylinder,data,degrees,del,% %
406     delete,depth,descender,diff_angle,digits,division,dot,draw_complete,% %
407     ellipsoid,emissive,end_face_color>equals,explod,extrusion,faces,factorial,% %
408     False,floor,follow,font,format,forward,fov,frame,gcurve,gdisplay,gdots,% %
409     get_library,get_selected,ghbars,global,GlowScript,graph,graphs,green,gvbars,% %
410     hat,headlength,headwidth,height,helix,hsv_to_rgb,index,interval,keydown,% %
411     keyup,label,length,lights,line,linecolor,linewidth,logx,logy,lower_left,% %
412     lower_right,mag,mag2,magenta,make_trail,marker_color,markers,material,% %
413     max,min,mouse,mousedown,mousemove,mouseup,newball,norm,normal,objects,% %
414     offset,one,opacity,orange,origin,path,pause,pi,pixel_to_world,pixels,plot,% %
415     points, pos, pow, pps, print, print_function, print_options, proj, purple, pyramid,% %
416     quad, radians, radius, random, rate, ray, read_local_file, readonly, red, redraw,% %
417     retain,rgb_to_hsv,ring,rotate,round,scene,scroll,shaftwidth,shape,shapes,% %
418     shininess,show_end_face,show_start_face,sign,sin,size,size_units,sleep,% %
419     smooth,space,sphere,sqrt,start,start_face_color,stop,tan,text,textpos,% %
420     texture,textures,thickness,title,trail_color,trail_object,trail_radius,% %
421     trail_type,triangle,trigger,True,twist,unbind,up,upper_left,upper_right,% %
422     userpan,userspin,userzoom,vec,vector,vertex,vertical_spacing,visible,% %
423     visual,vpython,VPython,waitfor,WebVPython,white,width,world,xtitle,% %
424     yellow,yoffset,ytitle% %
425   },%
426   morekeywords={print,None,TypeError},%           % additional keywords
427   morestring=[b]{""},%                         % treat triple quotes as strings
428   numbers=left,%                            % where to put line numbers
429   numbersep=10pt,%                          % how far line numbers are from code
430   numberstyle=\bf\tiny,%                      % set to 'none' for no line numbers
431   showstringspaces=false,%                  % show spaces in strings
432   showtabs=false,%                          % show tabs within strings
433   stringstyle=\sf\color{gsgreen},%             % color for strings
434   upquote=true,%                           % how to typeset quotes
435 }%

```

Introduce a new, more intelligent `\webpythonblock`<sup>P.62</sup> environment.

See <https://tex.stackexchange.com/a/232208/218142>.

```
436 \AtBeginEnvironment{webpythonblock}{\catcode`\#=12}
437 \AtEndEnvironment{webpythonblock}{\catcode`\#=6}
438 \NewTCBListing[auto counter,list inside=gsprogs]{webpythonblock}{ 0{} D(){webpython.org} m }%
439   {%
440     breakable,%
441     center,%
442     code = \newpage,%
443     %derivpeach,%
444     enhanced,%
445     hyperurl interior = https://#2,%
446     label = {gs:\thetcbcounter},%
447     left = 8mm,%
448     list entry = \thetcbcounter~~~~~#3,%
449     listing only,%
450     listing style = vpython,%
451     nameref = {#3},%
452     title = \begin{minipage}{1.5cm}%
453       \protect\qrcode*[https://#2]%
454       \end{minipage}\hspace{5mm}%
455     \begin{minipage}{0.7\textwidth}%
456       \texttt{Web VPython} Program \thetcbcounter: #3%
457     \end{minipage},%
458     width = 0.9\textwidth,%
459     {#1},%
460   }%
```

Here is a variant that omits the QR code.

```
461 \AtBeginEnvironment{webpythonblock*}{\catcode`\#=12}
462 \AtEndEnvironment{webpythonblock*}{\catcode`\#=6}
463 \NewTCBListing[use counter from=webpythonblock,list inside=gsprogs]
464   {webpythonblock*}{ 0{} D(){webpython.org} m }%
465   {%
466     breakable,%
467     center,%
468     code = \newpage,%
469     %derivpeach,%
470     enhanced,%
471     hyperurl interior = https://#2,%
472     label = {gs:\thetcbcounter},%
473     left = 8mm,%
474     list entry = \thetcbcounter~~~~~#3,%
475     listing only,%
476     listing style = vpython,%
477     nameref = {#3},%
478     title = \texttt{Web VPython} Program \thetcbcounter: #3,%
479     width = 0.9\textwidth,%
480     {#1},%
481   }%
```

A new command for generating a list of Web VPython programs.

```
482 \NewDocumentCommand{\listofwebpythonprograms}{}%
483   {%
484     \tcblistof[\section*]{gsprogs}{List of \texttt{Web VPython} Programs}%
485   }%
```

Introduce a new, more intelligent `\vpythonfile`<sup>P.68</sup> command.

See <https://tex.stackexchange.com/q/616205/218142>.

```

486 \newcommand{\vpythonfile}{\catcode`\#=12 \vpythonfile@auxA}
487 \NewDocumentCommand{\vpythonfile@auxA}{ O{} D() \vpython.org m m }%
488   {%
489     \vpythonfile@auxB[#1] (#2){#3}{#4}%
490     \catcode`\#=6
491   }%
492 \NewTCBInputListing[auto counter, list inside=vpprogs]
493   {\vpythonfile@auxB}{ O{} D() \vpython.org m m }%
494   {%
495     breakable,%
496     center,%
497     code = \newpage,%
498     %derivgray,%
499     enhanced,%
500     hyperurl interior = https://#2,%
501     label = {vp:\thetcbcounter},%
502     left = 8mm,%
503     list entry = \thetcbcounter~~~~~#4,%
504     listing file = {#3},%
505     listing only,%
506     listing style = vpython,%
507     nameref = {#4},%
508     title = \texttt{VPython} Program \thetcbcounter: #4,%
509     width = 0.9\textwidth,%
510     {#1},%
511   }%

```

A new command for generating a list of VPython programs.

```

512 \NewDocumentCommand{\listofvpythonprograms}{}%
513   {%
514     \tcblistof[\section*]{vpprogs}{List of \texttt{VPython} Programs}%
515   }%

```

Introduce a new `\webvpythoninline`<sup>P. 70</sup> command.

```

516 \DeclareTotalTCBox{\webvpythoninline}{ m }%
517   {%
518     bottom = Opt,%
519     bottomrule = 0.0mm,%
520     boxsep = 1.0mm,%
521     colback = gsbgrgray,%
522     colframe = gsbgrgray,%
523     left = Opt,%
524     leftrule = 0.0mm,%
525     nobeforeafter,%
526     right = Opt,%
527     rightrule = 0.0mm,%
528     sharp corners,%
529     tcbox raise base,%
530     top = Opt,%
531     toprule = 0.0mm,%
532   }%
533   {\lstinline[style = vpython]{#1}}%

```

Define `\vpythoninline`<sup>P. 70</sup>, a semantic alias for VPython in-line listings.

```

534 \NewDocumentCommand{\vpythoninline}{}{\webvpythoninline}%

```

## 6 The `mandiexp` Package

`mandi` comes with an accessory package `mandiexp` which includes commands specific to *Matter & Interactions*.<sup>5</sup> The commands are primarily for typesetting mathematical expressions used in that text. Note that `mandiexp` requires, and loads, `mandi` but `mandi` doesn't require, and doesn't load, `mandiexp`.

Load `mandiexp` as you would any package in your preamble. There are no package options.

```
\usepackage{mandiexp}
```

`\mandiexpversion`

Typesets the current version and build date.

```
The version is \mandiexpversion\ and is a stable build.
```

```
The version is v3.1.0 dated 2022-01-27 and is a stable build.
```

### 6.1 The Fundamental Principles

<code>\lhsmomentumprinciple</code>	(LHS of delta form, bold vectors)
<code>\rhsmomentumprinciple</code>	(RHS of delta form, bold vectors)
<code>\lhsmomentumprincipleupdate</code>	(LHS of update form, bold vectors)
<code>\rhsmomentumprincipleupdate</code>	(RHS of update form, bold vectors)
<code>\momentumprinciple</code>	(delta form, bold vectors)
<code>\momentumprincipleupdate</code>	(update form, bold vectors)
<code>\lhsmomentumprinciple*</code>	(LHS of delta form, arrow vectors)
<code>\rhsmomentumprinciple*</code>	(RHS of delta form, arrow vectors)
<code>\lhsmomentumprincipleupdate*</code>	(LHS of update form, arrow vectors)
<code>\rhsmomentumprincipleupdate*</code>	(RHS of update form, arrow vectors)
<code>\momentumprinciple*</code>	(delta form, arrow vectors)
<code>\momentumprincipleupdate*</code>	(update form, arrow vectors)

Variants of command for typesetting the momentum principle. Use starred variants to get arrow notation for vectors.

<sup>5</sup>See *Matter & Interactions* and <https://matterandinteractions.org/> for details.

$\backslash(\backslash\text{lhsmomentumprinciple} \backslash) \quad \backslash\backslash$	$\Delta p_{\text{sys}}$
$\backslash(\backslash\text{rhsmomentumprinciple} \backslash) \quad \backslash\backslash$	$\mathbf{F}_{\text{sys,net}} \Delta t$
$\backslash(\backslash\text{lhsmomentumprincipleupdate} \backslash) \quad \backslash\backslash$	$\mathbf{p}_{\text{sys,initial}}$
$\backslash(\backslash\text{rhsmomentumprincipleupdate} \backslash) \quad \backslash\backslash$	$\mathbf{p}_{\text{sys,initial}} + \mathbf{F}_{\text{sys,net}} \Delta t$
$\backslash(\backslash\text{momentumprinciple} \backslash) \quad \backslash\backslash$	$\Delta \mathbf{p}_{\text{sys}} = \mathbf{F}_{\text{sys,net}} \Delta t$
$\backslash(\backslash\text{momentumprincipleupdate} \backslash) \quad \backslash\backslash$	$\mathbf{p}_{\text{sys,final}} = \mathbf{p}_{\text{sys,initial}} + \mathbf{F}_{\text{sys,net}} \Delta t$
$\backslash(\backslash\text{lhsmomentumprinciple*} \backslash) \quad \backslash\backslash$	$\Delta \vec{p}_{\text{sys}}$
$\backslash(\backslash\text{rhsmomentumprinciple*} \backslash) \quad \backslash\backslash$	$\vec{\mathbf{F}}_{\text{sys,net}} \Delta t$
$\backslash(\backslash\text{lhsmomentumprincipleupdate*} \backslash) \quad \backslash\backslash$	$\vec{p}_{\text{sys,final}}$
$\backslash(\backslash\text{rhsmomentumprincipleupdate*} \backslash) \quad \backslash\backslash$	$\vec{p}_{\text{sys,initial}} + \vec{\mathbf{F}}_{\text{sys,net}} \Delta t$
$\backslash(\backslash\text{momentumprinciple*} \backslash) \quad \backslash\backslash$	$\Delta \vec{p}_{\text{sys}} = \vec{\mathbf{F}}_{\text{sys,net}} \Delta t$
$\backslash(\backslash\text{momentumprincipleupdate*} \backslash)$	$\vec{p}_{\text{sys,final}} = \vec{p}_{\text{sys,initial}} + \vec{\mathbf{F}}_{\text{sys,net}} \Delta t$

$\backslash\text{lhsenergyprinciple}$	(LHS of delta form)
$\backslash\text{rhsenergyprinciple}[(+process...)]$	(RHS of delta form)
$\backslash\text{lhsenergyprincipleupdate}$	(LHS of update form)
$\backslash\text{rhsenergyprincipleupdate}[(+process...)]$	(RHS of update form)
$\backslash\text{energyprinciple}[(+process...)]$	(delta form)
$\backslash\text{energyprincipleupdate}[(+process...)]$	(update form)

Variants of command for typesetting the energy principle.

$\backslash(\backslash\text{lhsenergyprinciple} \backslash) \quad \backslash\backslash$	$\Delta E_{\text{sys}}$
$\backslash(\backslash\text{rhsenergyprinciple} \backslash) \quad \backslash\backslash$	$W_{\text{ext}}$
$\backslash(\backslash\text{rhsenergyprinciple}[+Q] \backslash) \quad \backslash\backslash$	$W_{\text{ext}} + Q$
$\backslash(\backslash\text{energyprinciple} \backslash) \quad \backslash\backslash$	$\Delta E_{\text{sys}} = W_{\text{ext}}$
$\backslash(\backslash\text{energyprinciple}[+Q] \backslash) \quad \backslash\backslash$	$\Delta E_{\text{sys}} = W_{\text{ext}} + Q$
$\backslash(\backslash\text{lhsenergyprincipleupdate} \backslash) \quad \backslash\backslash$	$E_{\text{sys,final}}$
$\backslash(\backslash\text{rhsenergyprincipleupdate} \backslash) \quad \backslash\backslash$	$E_{\text{sys,initial}} + W_{\text{ext}}$
$\backslash(\backslash\text{rhsenergyprincipleupdate}[+Q] \backslash) \quad \backslash\backslash$	$E_{\text{sys,initial}} + W_{\text{ext}} + Q$
$\backslash(\backslash\text{energyprincipleupdate} \backslash) \quad \backslash\backslash$	$E_{\text{sys,final}} = E_{\text{sys,initial}} + W_{\text{ext}}$
$\backslash(\backslash\text{energyprincipleupdate}[+Q] \backslash)$	$E_{\text{sys,final}} = E_{\text{sys,initial}} + W_{\text{ext}} + Q$

$\backslash\text{lhsangularmomentumprinciple}$	(LHS of delta form, bold vectors)
$\backslash\text{rhsangularmomentumprinciple}$	(RHS of delta form, bold vectors)
$\backslash\text{lhsangularmomentumprincipleupdate}$	(LHS of update form, bold vectors)
$\backslash\text{rhsangularmomentumprincipleupdate}$	(RHS of update form, bold vectors)
$\backslash\text{angularmomentumprinciple}$	(delta form, bold vectors)
$\backslash\text{angularmomentumprincipleupdate}$	(update form, bold vectors)
$\backslash\text{lhsangularmomentumprinciple*}$	(LHS of delta form, arrow vectors)
$\backslash\text{rhsangularmomentumprinciple*}$	(RHS of delta form, arrow vectors)
$\backslash\text{lhsangularmomentumprincipleupdate*}$	(LHS of update form, arrow vectors)
$\backslash\text{rhsangularmomentumprincipleupdate*}$	(RHS of update form, arrow vectors)
$\backslash\text{angularmomentumprinciple*}$	(delta form, arrow vectors)

### \angularmomentumprincipleupdate\*

(update form, arrow vectors)

Variants of command for typesetting the angular momentum principle. Use starred variants to get arrow notation for vectors.

```
\(\lhsangularmomentumprinciple \)    \\
\(\rhsangularmomentumprinciple \)    \\
\(\lhsangularmomentumprincipleupdate \) \\
\(\rhsangularmomentumprincipleupdate \) \\
\(\angularmomentumprinciple \)        \\
\(\angularmomentumprincipleupdate \)   \\
\(\lhsangularmomentumprinciple* \)   \\
\(\rhsangularmomentumprinciple* \)   \\
\(\lhsangularmomentumprincipleupdate* \) \\
\(\rhsangularmomentumprincipleupdate* \) \\
\(\angularmomentumprinciple* \)      \\
\(\angularmomentumprincipleupdate* \)
```

$$\begin{aligned} & \Delta \mathbf{L}_{A,\text{sys},\text{net}} \\ & \tau_{A,\text{sys},\text{net}} \Delta t \\ & \mathbf{L}_{A,\text{sys},\text{final}} \\ & \mathbf{L}_{A,\text{sys},\text{initial}} + \tau_{A,\text{sys},\text{net}} \Delta t \\ & \Delta \vec{\mathbf{L}}_{A,\text{sys},\text{net}} = \tau_{A,\text{sys},\text{net}} \Delta t \\ & \mathbf{L}_{A,\text{sys},\text{final}} = \mathbf{L}_{A,\text{sys},\text{initial}} + \tau_{A,\text{sys},\text{net}} \Delta t \\ & \Delta \vec{\mathbf{L}}_{A,\text{sys},\text{net}} \\ & \vec{\tau}_{A,\text{sys},\text{net}} \Delta t \\ & \vec{\mathbf{L}}_{A,\text{sys},\text{final}} \\ & \vec{\mathbf{L}}_{A,\text{sys},\text{initial}} + \vec{\tau}_{A,\text{sys},\text{net}} \Delta t \\ & \Delta \vec{\mathbf{L}}_{A,\text{sys},\text{net}} = \vec{\tau}_{A,\text{sys},\text{net}} \Delta t \\ & \vec{\mathbf{L}}_{A,\text{sys},\text{final}} = \vec{\mathbf{L}}_{A,\text{sys},\text{initial}} + \vec{\tau}_{A,\text{sys},\text{net}} \Delta t \end{aligned}$$

## 6.2 Other Expressions

N 2021-02-13

### \energyof{<label>} [<label>]

Generic symbol for the energy of some entity.

```
\(\energyof{\symup{electron}} \) \\
\(\energyof{\symup{electron}}[\symup{final}] \)
```

$$\begin{aligned} & E_{\text{electron}} \\ & E_{\text{electron},\text{final}} \end{aligned}$$

N 2021-02-13

### \systemenergy [<label>]

Symbol for system energy.

```
\(\systemenergy \) \\
\(\systemenergy[\symup{final}] \)
```

$$\begin{aligned} & E_{\text{sys}} \\ & E_{\text{sys},\text{final}} \end{aligned}$$

N 2021-02-13

### \particleenergy [<label>]

Symbol for particle energy.

```
\(\particleenergy \) \\
\(\particleenergy[\symup{final}] \)
```

$$\begin{aligned} & E_{\text{particle}} \\ & E_{\text{particle},\text{final}} \end{aligned}$$

N 2021-02-13

### \restenergy [<label>]

Symbol for rest energy.

```
\(\restenergy\)\  
\(\restenergy[\symup{final}]\)\
```

$E_{\text{rest}}$   
 $E_{\text{rest,final}}$

#### N 2021-02-13 **\internalenergy** [*<label>*]

Symbol for internal energy.

```
\(\internalenergy\)\  
\(\internalenergy[\symup{final}]\)\
```

$E_{\text{internal}}$   
 $E_{\text{internal,final}}$

#### N 2021-02-13 **\chemicalenergy** [*<label>*]

Symbol for chemical energy.

```
\(\chemicalenergy\)\  
\(\chemicalenergy[\symup{final}]\)\
```

$E_{\text{chem}}$   
 $E_{\text{chem,final}}$

#### N 2021-02-13 **\thermalenergy** [*<label>*]

Symbol for thermal energy.

```
\(\thermalenergy\)\  
\(\thermalenergy[\symup{final}]\)\
```

$E_{\text{therm}}$   
 $E_{\text{therm,final}}$

#### N 2021-02-13 **\photonenergy** [*<label>*]

Symbol for photon energy.

```
\(\photonenergy\)\  
\(\photonenergy[\symup{final}]\)\
```

$E_{\text{photon}}$   
 $E_{\text{photon,final}}$

#### N 2021-02-13 **\translationalkineticenergy** [*<label>*]

#### N 2021-02-13 **\translationalkineticenergy\*** [*<label>*]

Symbol for translational kinetic energy. The starred variant gives  $E$  notation.

```
\(\translationalkineticenergy\)\  
\(\translationalkineticenergy[\symup{initial}]\)\  
\(\translationalkineticenergy*\)\  
\(\translationalkineticenergy*[\symup{initial}]\)\
```

$K_{\text{trans}}$   
 $K_{\text{trans,initial}}$   
 $E_K$   
 $E_{K,\text{initial}}$

N 2021-02-13  
N 2021-02-13

\rotationalkineticenergy[*label*]  
\rotationalkineticenergy\*[*label*]

Symbol for rotational kinetic energy. The starred variant gives  $E$  notation.

```
\(\ \rotationalkineticenergy \)      \\
\(\ \rotationalkineticenergy[\symup{initial}] \) \\
\(\ \rotationalkineticenergy* \)      \\
\(\ \rotationalkineticenergy*[\symup{initial}] \)
```

$K_{\text{rot}}$   
 $K_{\text{rot,initial}}$   
 $E_{\text{rot}}$   
 $E_{\text{rot,initial}}$

N 2021-02-13  
N 2021-02-13

\vibrationalkineticenergy[*label*]  
\vibrationalkineticenergy\*[*label*]

Symbol for vibrational kinetic energy. The starred variant gives  $E$  notation.

```
\(\ \vibrationalkineticenergy \)      \\
\(\ \vibrationalkineticenergy[\symup{initial}] \) \\
\(\ \vibrationalkineticenergy* \)      \\
\(\ \vibrationalkineticenergy*[\symup{initial}] \)
```

$K_{\text{vib}}$   
 $K_{\text{vib,initial}}$   
 $E_{\text{vib}}$   
 $E_{\text{vib,initial}}$

N 2021-02-13

\gravitationalpotentialenergy[*label*]

Symbol for gravitational potential energy.

```
\(\ \gravitationalpotentialenergy \) \\
\(\ \gravitationalpotentialenergy[\symup{final}] \)
```

$U_g$   
 $U_{g,\text{final}}$

N 2021-02-13

\electricpotentialenergy[*label*]

Symbol for electric potential energy.

```
\(\ \electricpotentialenergy \) \\
\(\ \electricpotentialenergy[\symup{final}] \)
```

$U_e$   
 $U_{e,\text{final}}$

N 2021-02-13

\springpotentialenergy[*label*]

Symbol for spring potential energy.

```
\(\ \springpotentialenergy \) \\
\(\ \springpotentialenergy[\symup{final}] \)
```

$U_s$   
 $U_{s,\text{final}}$

### 6.3 **mandiexp** Source Code

Definine the package version and date for global use, exploiting the fact that in a .sty file there is now no need for `\makeatletter` and `\makeatother`. This simplifies defining internal commands, with @ in the name, that are not for the user to know about.

```
1 \def\mandiexp@version{3.1.0}
2 \def\mandiexp@date{2022-01-27}
3 \NeedsTeXFormat{LaTeX2e}[2020-02-02]
4 \DeclareRelease{v3.1.0}{2022-01-27}{\mandiexp.sty}
5 \DeclareCurrentRelease{v\mandiexp@version}{\mandiexp@date}
6 \ProvidesPackage{mandiexp}
7   [\mandiexp@date\space v\mandiexp@version\space Macros for Matter & Interactions]

Define a convenient package version command.

8 \newcommand*\mandiexpversion{v\mandiexp@version\space dated \mandiexp@date}
9 \RequirePackage{mandi}

10 \IfFormatAtLeastTF {2020-10-01} % load xparse if necessary
11   {}%
12   {\RequirePackage{xparse}}%

13 \typeout{}%
14 \typeout{\mandiexp: You are using mandiexp \mandiexpversion.}%
15 \typeout{\mandiexp: This package requires LuaLaTeX.}%
16 \typeout{}%
17 %
18 % Commands specific to Matter & Interactions
19 % The momentum principle
20 \NewDocumentCommand{\lhsmomentumprinciple}{ s }%
21   {%
22     \Delta
23     \IfBooleanTF{#1}%
24       {%
25         \vec*{p}%
26       }%
27       {%
28         \vec{p}%
29       }%
30     \sb{\symup{sys}}%
31   }%
32 \NewDocumentCommand{\rhsmomentumprinciple}{ s }%
33   {%
34     \IfBooleanTF{#1}%
35       {%
36         \vec*{F}%
37       }%
38       {%
39         \vec{F}%
40       }%
41     \sb{\symup{sys,net}}\,,\Delta t%
42   }%
43 \NewDocumentCommand{\lhsmomentumprincipleupdate}{ s }%
44   {%
45     \IfBooleanTF{#1}%
46       {%
47         \vec*{p}%
48       }%
49       {%
50         \vec{p}%
51       }%
```

```

52     \sb{\symup{sys,final}}%
53   }%
54 \NewDocumentCommand{\rhsmomentumprincipleupdate}{ s }%
55   {%
56     \IfBooleanTF{#1}%
57       {%
58         \vec*{p}%
59       }%
60       {%
61         \vec{p}%
62       }%
63     \sb{\symup{sys,initial}}+%
64     \IfBooleanTF{#1}%
65       {%
66         \vec*{F}%
67       }%
68       {%
69         \vec{F}%
70       }%
71     \sb{\symup{sys,net}}\Delta t%
72   }%
73 \NewDocumentCommand{\momentumprinciple}{ s }%
74   {%
75     \IfBooleanTF{#1}%
76       {%
77         \lhsmomentumprinciple* = \rhsmomentumprinciple*%
78       }%
79       {%
80         \lhsmomentumprinciple = \rhsmomentumprinciple%
81       }%
82   }%
83 \NewDocumentCommand{\momentumprincipleupdate}{ s }%
84   {%
85     \IfBooleanTF{#1}%
86       {%
87         \lhsmomentumprincipleupdate* = \rhsmomentumprincipleupdate*%
88       }%
89       {%
90         \lhsmomentumprincipleupdate = \rhsmomentumprincipleupdate%
91       }%
92   }%
93 % The momentum principle
94 \NewDocumentCommand{\lhsenergyprinciple}{}%
95   {%
96     \Delta E_{\symup{sys}}%
97   }%
98 \NewDocumentCommand{\rhsenergyprinciple}{ 0{} }%
99   {%
100    W_{\symup{ext}}#1%
101  }%
102 \NewDocumentCommand{\lhsenergyprincipleupdate}{}%
103   {%
104     E_{\symup{sys,final}}%
105   }%
106 \NewDocumentCommand{\rhsenergyprincipleupdate}{ 0{} }%
107   {%
108     E_{\symup{sys,initial}}+%
109     W_{\symup{ext}}#1%
110   }%

```

```

111 \NewDocumentCommand{\energyprinciple}{ O{} }%
112   {%
113     \lhsenergyprinciple = \rhsenergyprinciple[#1]%
114   }%
115 \NewDocumentCommand{\energyprincipleupdate}{ O{} }%
116   {%
117     \lhsenergyprincipleupdate = \rhsenergyprincipleupdate[#1]%
118   }%
119 % The angular momentum principle
120 \NewDocumentCommand{\lhsangularmomentumprinciple}{ s }%
121   {%
122     \Delta%
123     \IfBooleanTF{#1}%
124       {%
125         \vec*{L}%
126       }%
127       {%
128         \vec{L}%
129       }%
130     \sb{A\symup{,sys,net}}%
131   }%
132 \NewDocumentCommand{\rhsangularmomentumprinciple}{ s }%
133   {%
134     \IfBooleanTF{#1}%
135       {%
136         \vec*{\tau}%
137       }%
138       {%
139         \vec{\tau}%
140       }%
141     \sb{A\symup{,sys,net}}\,,\Delta t%
142   }%
143 \NewDocumentCommand{\lhsangularmomentumprincipleupdate}{ s }%
144   {%
145     \IfBooleanTF{#1}%
146       {%
147         \vec*{L}%
148       }%
149       {%
150         \vec{L}%
151       }%
152     \sb{A,\symup{sys,final}}%
153   }%
154 \NewDocumentCommand{\rhsangularmomentumprincipleupdate}{ s }%
155   {%
156     \IfBooleanTF{#1}%
157       {%
158         \vec*{L}%
159       }%
160       {%
161         \vec{L}%
162       }%
163     \sb{A\symup{,sys,initial}}+%
164     \IfBooleanTF{#1}%
165       {%
166         \vec*{\tau}%
167       }%
168       {%
169         \vec{\tau}%

```

```

170      }%
171      \sb{A\symup{,sys,net}}\,\Delta t%
172  }%
173 \NewDocumentCommand{\angularmomentumprinciple}{ s }%
174 {%
175   \IfBooleanTF{#1}%
176   {%
177     \lhsangularmomentumprinciple* = \rhsangularmomentumprinciple*%
178   }%
179   {%
180     \lhsangularmomentumprinciple = \rhsangularmomentumprinciple%
181   }%
182 }%
183 \NewDocumentCommand{\angularmomentumprincipleupdate}{ s }%
184 {%
185   \IfBooleanTF{#1}%
186   {%
187     \lhsangularmomentumprincipleupdate* = \rhsangularmomentumprincipleupdate*%
188   }%
189   {%
190     \lhsangularmomentumprincipleupdate = \rhsangularmomentumprincipleupdate%
191   }%
192 }%
193 \NewDocumentCommand{\energyof}{ m o }%
194 {%
195   E_{#1}%
196   \IfValueT{#2}%
197   {, #2}%
198 }%
199 }%
200 \NewDocumentCommand{\systemenergy}{ o }%
201 {%
202   E_{\symup{sys}}%
203   \IfValueT{#1}%
204   {, #1}%
205 }%
206 }%
207 \NewDocumentCommand{\particleenergy}{ o }%
208 {%
209   E_{\symup{particle}}%
210   \IfValueT{#1}%
211   {, #1}%
212 }%
213 }%
214 \NewDocumentCommand{\restenergy}{ o }%
215 {%
216   E_{\symup{rest}}%
217   \IfValueT{#1}%
218   {, #1}%
219 }%
220 }%
221 \NewDocumentCommand{\internalenergy}{ o }%
222 {%
223   E_{\symup{internal}}%
224   \IfValueT{#1}%
225   {, #1}%
226 }%
227 }%
228 \NewDocumentCommand{\chemicalenergy}{ o }%

```

```

229   {%
230     E_{\symup{chem}}%
231     \IfValueT{#1}%
232     {,#1}%
233   }%
234 }%
235 \NewDocumentCommand{\thermalenergy}{ o }{%
236   {%
237     E_{\symup{therm}}%
238     \IfValueT{#1}%
239     {,#1}%
240   }%
241 }%
242 \NewDocumentCommand{\photonenergy}{ o }{%
243   {%
244     E_{\symup{photon}}%
245     \IfValueT{#1}%
246     {,#1}%
247   }%
248 }%
249 \NewDocumentCommand{\translationalkineticenergy}{ s d[] }{%
250   {%
251     % d[] must be used because of the way consecutive optional
252     % arguments are handled. See xparse docs for details.
253     % See https://tex.stackexchange.com/a/569011/218142
254     \IfBooleanTF{#1}%
255     {%
256       E_{\bgroup \symup{K}}%
257     }%
258     {%
259       K_{\bgroup\symup{trans}}%
260     }%
261     \IfValueT{#2}{, #2}%
262     \egroup%
263   }%
264 \NewDocumentCommand{\rotationalkineticenergy}{ s d[] }{%
265   {%
266     % d[] must be used because of the way consecutive optional
267     % arguments are handled. See xparse docs for details.
268     % See https://tex.stackexchange.com/a/569011/218142
269     \IfBooleanTF{#1}%
270     {%
271       E_{\bgroup}%
272     }%
273     {%
274       K_{\bgroup}%
275     }%
276     \symup{rot}\IfValueT{#2}{, #2}%
277     \egroup%
278   }%
279 \NewDocumentCommand{\vibrationalkineticenergy}{ s d[] }{%
280   {%
281     % d[] must be used because of the way consecutive optional
282     % arguments are handled. See xparse docs for details.
283     % See https://tex.stackexchange.com/a/569011/218142
284     \IfBooleanTF{#1}%
285     {%
286       E_{\bgroup}%
287     }%

```

```

288      {%
289      K_ \bgroup%
290      }%
291      \symup{vib}\IfValueT{#2}{, #2}%
292      \egroup%
293      }%
294 \NewDocumentCommand{\gravitationalpotentialenergy}{ o }{%
295      {%
296      U_{\symup{g}}%
297      \IfValueT{#1}{%
298          {, #1}%
299      }%
300      }%
301 \NewDocumentCommand{\electricpotentialenergy}{ o }{%
302      {%
303      U_{\symup{e}}%
304      \IfValueT{#1}{%
305          {, #1}%
306      }%
307      }%
308 \NewDocumentCommand{\springpotentialenergy}{ o }{%
309      {%
310      U_{\symup{s}}%
311      \IfValueT{#1}{%
312          {, #1}%
313      }%
314      }%

```

## 7 Index

Page numbers refer to page where the corresponding entry is documented and/or referenced.

### A

\absolutevalue	54
\absolutevalue*	54
\acceleration	10
\accelerationvector	10
alternate value	8, 36
\alwaysusealternateunits	22
\alwaysuseapproximateconstants	31
\alwaysusebaseunits	22
\alwaysusederivedunits	22
\alwaysusepreciseconstants	31
\amount	10
\ampere	31
\anglebrackets	52
\anglebrackets*	52
\angularacceleration	10
\angularaccelerationvector	10
\angularfrequency	10
\angularimpulse	10
\angularimpulsevector	10
\angularmomentum	11
\angularmomentumprinciple	83
\angularmomentumprinciple*	83
\angularmomentumprincipleupdate	83
\angularmomentumprincipleupdate*	84
\angularmomentumvector	11
\angularvelocity	11
\angularvelocityvector	11
\area	11
\areachargedensity	11
\areamassdensity	11
\atomicmassunit	32
\avogadro	24

### B

base value	8, 36
\biotsavartconstant	24
\bohrradius	24
\boltzmann	25

### C

\candela	32
\capacitance	11
\changein	52
\charge	11
\checkconstant	24

\checkquantity	10
\chemicalenergy	85
\cmagneticfield	12
\cmagneticfieldvector	12
\colvec	60
\conductance	12
\conductivity	12
\contraction	61
\contraction*	61
\conventionalcurrent	12
\coulomb	32
\coulombconstant	25
\curlybraces	53
\curlybraces*	53
\current	12
\currentdensity	12
\currentdensityvector	12

### D

\degree	32
derived value	8, 36
\df	61
\dielectricconstant	12
\direction	13
\directionvector	13
\dirvec	52
\dirvec*	52
\displacement	13
\displacementvector	13
\doublebars	52
\doublebars*	52
\duration	13

### E

\earthmass	25
\earthmoondistance	25
\earthradius	25
\earthsundistance	25
\electricdipolemoment	13
\electricdipolemomentvector	13
\electricfield	13
\electricfieldvector	13
\electricflux	13
\electricpotential	13
\electricpotentialdifference	14
\electricpotentialenergy	86

\electronCharge	26	\henry	32
\electroncharge	26	\hereuseapproximateconstants	31
\electroncurrent	14	\hereusebaseunits	22
\electronmass	26	\hereusedalternateunits	22
\electronvolt	32	\hereusedderivedunits	22
\elementarycharge	26	\hereusepreciseconstants	31
\emf	14	\hertz	32
\emptyunit	31	\hilite	58
\energy	14	\hydrogenmass	26
\energydensity	14		
\energyflux	15		
\energyfluxvector	15		
\energyinev	14		
\energyinkev	14		
\energyinmev	14		
\energyof	84		
\energyprinciple	83		
\energyprincipleupdate	83		
\entropy	15		
Environments			
parts	55		
physicsproblem	55		
physicsproblem*	55		
physicssolution	56		
physicssolution*	56		
usealternateunits	23		
useapproximateconstants	31		
usebaseunits	23		
usederivedunits	23		
usepreciseconstants	31		
webpythonblock	62		
webpythonblock*	62		
\ev	32		
F			
false value	8, 36		
\farad	32		
\finestructure	26		
\force	15		
\forcevector	15		
\frequency	15		
G			
\gravitationalfield	15		
\gravitationalfieldvector	15		
\gravitationalpotential	15		
\gravitationalpotentialdifference	15		
\gravitationalpotentialenergy	86		
H			
\hbar	33		
I			
\image	59		
\impulse	16		
\impulsevector	16		
\indexofrefraction	16		
\inductance	16		
\internalenergy	85		
\inverse	32		
J			
\joule	32		
K			
\kelvin	32		
\kev	32		
Keys			
preciseconstants	8		
units	8		
\kilolectronvolt	32		
\kilogram	32		
L			
\lhsangularmomentumprinciple	83		
\lhsangularmomentumprinciple*	83		
\lhsangularmomentumprincipleupdate	83		
\lhsangularmomentumprincipleupdate*	83		
\lhsenergyprinciple	83		
\lhsenergyprincipleupdate	83		
\lhsmomentumprinciple	82		
\lhsmomentumprinciple*	82		
\lhsmomentumprincipleupdate	82		
\lhsmomentumprincipleupdate*	82		
\lightspeed	32		
\linearchargedensity	16		
\linearmassdensity	16		
\lorentzfactor	16		
\luminousintensity	16		
M			
\magneticcharge	17		
\magneticdipolemoment	17		

\magneticdipolemomentvector . . . . .	17	\particleenergy . . . . .	84
\magneticfield . . . . .	17	parts environment . . . . .	55
\magneticfieldvector . . . . .	17	\pascal . . . . .	32
\magneticflux . . . . .	17	\per . . . . .	31
\magnitude . . . . .	54	\permeability . . . . .	18
\magnitude* . . . . .	54	\permittivity . . . . .	18
\mandiexpversion . . . . .	82	\perpendicularto . . . . .	55
\mandisetup . . . . .	8	\photonenergy . . . . .	85
\mandistudentversion . . . . .	51	physicsproblem environment . . . . .	55
\mandiversion . . . . .	8	physicsproblem* environment . . . . .	55
\mass . . . . .	17	physicssolution environment . . . . .	56
\megaelectronvolt . . . . .	32	physicssolution* environment . . . . .	56
\meter . . . . .	32	\planck . . . . .	28
\metre . . . . .	32	\planckbar . . . . .	28
\mev . . . . .	32	\planckc . . . . .	28
\mivector . . . . .	34	\planeangle . . . . .	18
\mobility . . . . .	17	\polarizability . . . . .	18
\mole . . . . .	32	\power . . . . .	18
\momentofinertia . . . . .	17	\poynting . . . . .	19
\momentum . . . . .	9, 18	\poyntingvector . . . . .	19
\momentumflux . . . . .	18	preciseconstants key . . . . .	8
\momentumfluxvector . . . . .	18	\pressure . . . . .	19
\momentumprinciple . . . . .	82	\problempart . . . . .	55
\momentumprinciple* . . . . .	82	\protonCharge . . . . .	28
\momentumprincipleupdate . . . . .	82	\protoncharge . . . . .	28
\momentumprincipleupdate* . . . . .	82	\protonmass . . . . .	29
\momentumvector . . . . .	9, 18		
\moonearthdistance . . . . .	27		
\moonmass . . . . .	27		
\moonradius . . . . .	27		
\mzofp . . . . .	27		
		<b>R</b>	
		\radian . . . . .	32
		\reason . . . . .	57
		\relativepermeability . . . . .	19
		\relativepermittivity . . . . .	19
		\renewphysicalconstant . . . . .	30
		\renewscalarquantity . . . . .	22
		\renewvectorquantity . . . . .	22
		\resistance . . . . .	19
		\resistivity . . . . .	19
		\restenergy . . . . .	84
		\rhsangularmomentumprinciple . . . . .	83
		\rhsangularmomentumprinciple* . . . . .	83
		\rhsangularmomentumprincipleupdate . . . . .	83
		\rhsangularmomentumprincipleupdate* . . . . .	83
		\rhsenergyprinciple . . . . .	83
		\rhsenergyprincipleupdate . . . . .	83
		\rhsmomentumprinciple . . . . .	82
		\rhsmomentumprinciple* . . . . .	82
		\rhsmomentumprincipleupdate . . . . .	82
		\rhsmomentumprincipleupdate* . . . . .	82
		\rotationalkineticenergy . . . . .	86
		\rotationalkineticenergy* . . . . .	86

\rowvec	60
\rydberg	29
<b>S</b>	
\second	32
\siemens	32
\singlebars	52
\singlebars*	52
\slot	61
\slot*	61
\solidangle	19
\specificheatcapacity	20
\speedoflight	29
\springpotentialenergy	86
\springstiffness	20
\springstretch	20
\squarebrackets	53
\squarebrackets*	53
\stefanboltzmann	29
\steradian	32
\strain	20
\stress	20
\sunearthdistance	29
\sunradius	29
\surfacegravfield	30
\systemenergy	84
<b>T</b>	
\temperature	20
\tencomp	60
\tencomp*	60
\tento	34
\tesla	32
\thermalenergy	85
\timestento	34
\torque	20
\torquevector	20
\tothefour	32
\totheinversefour	32
\totheinversethree	32
\totheinversetwo	32
\tothethree	32
\tothetwo	32
\translationalkineticenergy	85
\translationalkineticenergy*	85
true value	8, 36
<b>U</b>	
\unit	31
units key	8
\universalgrav	30
usealternateunits environment	23
useapproximateconstants environment	31
usebaseunits environment	23
usederivedunits environment	23
usepreciseconstants environment	31
\usk	31
<b>V</b>	
\vacuumpermeability	30
\vacuumpermittivity	30
\valence	61
\valence*	61
Values	
alternate	8, 36
base	8, 36
derived	8, 36
false	8, 36
true	8, 36
\vec	51
\vec*	51
\veccomp	60
\veccomp*	60
\vectoracceleration	10
\vectorangularacceleration	10
\vectorangularimpulse	10
\vectorangularmomentum	11
\vectorangularvelocity	11
\vectorcmagneticfield	12
\vectorcurrentdensity	12
\vectordirection	13
\vectordisplacement	13
\vectorelectricdipolemoment	13
\vectorelectricfield	13
\vectorenergyflux	15
\vectorforce	15
\vectorgravitationalfield	15
\vectorimpulse	16
\vectormagneticdipolemoment	17
\vectormagneticfield	17
\vectormomentum	9, 18
\vectormomentumflux	18
\vectorpoynting	19
\vectortorque	20
\vectorvelocity	20
\vectorvelocityc	21
\vectorwavenumber	21
\velocity	20
\velocityc	21
\velocitycvector	21
\velocityvector	20

\vibrationalkineticenergy	86
\vibrationalkineticenergy*	86
\volt	32
\volume	21
\volumechargedensity	21
\volumemassdensity	21
\vpythonfile	68
\vpythoninline	70

## W

\watt	32
\wavelength	21
\wavenumber	21
\wavenumbervector	21
\weber	32
webpythonblock environment	62
webpythonblock* environment	62
\webpythoninline	70
\work	21

## X

\xtento	34
---------	----

## Y

\youngsmodulus	22
----------------	----

## Z

\zerovec	52
\zerovec*	52